

Cancer Stem Cells: Models and Concepts

Piero Dalerba, Robert W. Cho,
and Michael F. Clarke

Stanford Institute for Stem Cell Biology and Regenerative Medicine, Stanford
University, Stanford, California 94304; email: pdalerba@stanford.edu,
robcho@stanford.edu, mclarke@stanford.edu

Annu. Rev. Med. 2007. 58:267–84

First published online as a Review in Advance on
September 26, 2006

The *Annual Review of Medicine* is online at
<http://med.annualreviews.org>

This article's doi:
10.1146/annurev.med.58.062105.204854

Copyright © 2007 by Annual Reviews.
All rights reserved

0066-4219/07/0218-0267\$20.00

Key Words

self-renewal, aberrant differentiation, functional heterogeneity of
tumor tissues, targeting of antitumor treatments

Abstract

Although monoclonal in origin, most tumors appear to contain a heterogeneous population of cancer cells. This observation is traditionally explained by postulating variations in tumor microenvironment and coexistence of multiple genetic subclones, created by progressive and divergent accumulation of independent somatic mutations. An additional explanation, however, envisages human tumors not as mere monoclonal expansions of transformed cells, but rather as complex tridimensional tissues where cancer cells become functionally heterogeneous as a result of differentiation. According to this second scenario, tumors act as caricatures of their corresponding normal tissues and are sustained in their growth by a pathological counterpart of normal adult stem cells, cancer stem cells. This model, first developed in human myeloid leukemias, is today being extended to solid tumors, such as breast and brain cancer. We review the biological basis and the therapeutic implications of the stem cell model of cancer.

DEFINITION OF “CANCER STEM CELLS”

Many human tissues undergo rapid and continuous cell turnover. In the colonic mucosa or in the peripheral blood, for example, the average life span of a mature, differentiated cell (e.g., a goblet cell in a crypt of the large intestine or a circulating granulocyte) can be measured in days or even hours. Despite the ephemeral nature of most of their individual cell components, human tissues maintain their mass and architecture over time through a tightly regulated process of renovation. Under physiological conditions, this process is sustained by a small minority of long-lived cells with extraordinary expansion potential, known as stem cells. Stem cells are defined by three main properties:

1. differentiation—the ability to give rise to a heterogeneous progeny of cells, which progressively diversify and specialize according to a hierarchical process, constantly replenishing the tissue of short-lived, mature elements;
2. self-renewal—the ability to form new stem cells with identical, intact potential for proliferation, expansion, and differentiation, thus maintaining the stem cell pool;
3. homeostatic control—the ability to modulate and balance differentiation and self-renewal according to environmental stimuli and genetic constraints.

Like their normal tissue counterparts, tumors are composed of heterogeneous populations of cells that differ in their apparent state of differentiation. Indeed, the differentiation features of a tumor, morphological and architectural, are the key parameter used in routine clinical practice by surgical pathologists to define a tumor's primary anatomical origin. This simple observation suggests that tumors are not mere monoclonal expansions of cells but might actually be akin to “abnormal organs,” sustained by a diseased “cancer stem cell” (CSC) population, which is endowed with the ability to self-renew and

undergo aberrant differentiation (1, 2). This hypothesis is further reinforced by the fact that cancer is known to result from the accumulation of multiple genetic mutations in a single target cell, sometimes over a period of many years (3). Because stem cells are the only long-lived cells in many tissues, they are the natural candidates in which early transforming mutations may accumulate.

Our limited knowledge of normal stem cells, in part due to the overall paucity of experimental assays for their functional study, has made the CSC theory difficult to probe. A new wave of studies, however, has recently begun to address this concept using an innovative, purely empirical approach, based on an *in vivo* self-renewal assay (4). Starting from whole tumor tissues, cancer cells are purified into single-cell suspensions and subsequently fractionated in different subsets according to the expression of a specific repertoire of surface markers. Once isolated, individual cancer cell subsets are injected into appropriate hosts (in most cases orthotopic tissues of immunodeficient mouse strains), and the subsets are compared with respect to tumorigenic capacity. According to the CSC model, only a specific subset of the cancer cell population (i.e., the long-lived CSC subset) should be able to sustain *in vivo* tumor growth, whereas all other subsets (i.e., the tumor counterparts of short-lived differentiated cells) should not. Indeed, this assumption has now been repeatedly confirmed in several tumor systems. Three key observations classically define the existence of a CSC population:

1. Only a minority of cancer cells within each tumor are usually endowed with tumorigenic potential when transplanted into immunodeficient mice.
2. Tumorigenic cancer cells are characterized by a distinctive profile of surface markers and can be differentially and reproducibly isolated from nontumorigenic ones by means of flow cytometry or other immunoselection procedures.
3. Tumors grown from tumorigenic cells contain mixed populations of

tumorigenic and nontumorigenic cancer cells, thus recreating the full phenotypic heterogeneity of the parent tumor.

It is important to note that, based on this approach, the term cancer stem cells represents a working definition with a purely operational significance. The term is used to indicate a tumor-initiating cell subset that can give rise to a heterogeneous progeny, similar in composition to the tissue from which it was originally isolated. In most cases, it is currently not possible to define with certainty the “genealogical” relationship between CSCs and normal stem cells of the corresponding tissues (i.e., whether CSCs originate directly from normal stem cells or the early stages of their progeny). Irrespective of the actual origins of CSCs, the identification of a CSC population establishes a functional hierarchy within a tumor tissue and encompasses both the self-renewal and differentiation hallmarks of stem cells. First developed in human myeloid leukemias, the CSC working model is today being progressively extended to several solid tumors, along with several biological and therapeutic implications.

DEVELOPMENT OF THE CANCER STEM CELL MODEL IN HUMAN MYELOID LEUKEMIAS

Normal Hematopoietic Stem Cells

Among mammalian tissues, the hematopoietic system is the first- and best-characterized in terms of hierarchical organization and sequential differentiation of cellular subpopulations. Starting in the 1960s, transplantation experiments in mice demonstrated the existence in the bone marrow of clonogenic precursors capable of remarkable long-term expansion and multipotent myelo-erythroid differentiation (5). These studies allowed progressive functional and phenotypic dissection of blood's distinct cellular lineages and lineage precursors (6). They provided most of the conceptual framework and the technical

terminology for the interpretation of adult stem cell biology. They were founded on two main observations:

1. In lethally irradiated mice, where native bone marrow has been completely eradicated, all the different hematopoietic lineages can be fully and permanently reconstituted by transplantation of a very small population of cells, representing as little as 0.05% of total bone marrow. This cell population is characterized by a specific surface marker phenotype that, remarkably, is negative for expression of all lineage-specific differentiation antigens (Lin^{neg}) (7).
2. Cells capable of multi-lineage reconstitution can be further subdivided into two distinct functional groups: long-term reconstituting multipotent progenitors, also defined as true hematopoietic stem cells (HSCs), and short-term or transiently reconstituting multipotent progenitors (MPPs). Both HSCs and MPPs can sustain the production of the full spectrum of blood cells, but reconstitution driven by HSCs is permanent and can last for the entire life of the transplanted animal, whereas reconstitution driven by MPPs is temporary and is usually exhausted after 2–3 months (8). Most importantly, HSCs and MPPs are organized according to a hierarchical lineage: HSCs give rise to MPPs (9).

Taken together, these observations indicate that the long-term, continuous renovation of a complex, functionally heterogeneous tissue (blood) can be sustained by a tiny, undifferentiated population of cells (HSCs), which is capable of both self-renewal and differentiation into a numerically large progeny. According to this model, the differentiation process takes place in two steps: first the loss of self-renewal capacity (the differentiation of HSCs into MPPs), then the actual progressive differentiation into the full spectrum of blood cells, which involves multiple stages and




Figure 1

Models of normal and malignant hematopoiesis. (*a*) Blood cells originate from hematopoietic stem cells (HSCs). HSCs differentiate into multipotent progenitors (MPPs), which in turn give rise to oligo-lineage precursors, such as the common lymphoid progenitor (CLP) and common myeloid progenitor (CMP). CLPs differentiate into T, B, and NK cells, whereas CMPs differentiate into macrophages (MF), neutrophils (PMN), erythrocytes (E), and platelets (PLT). (*b*) In humans, the early stages of hematopoiesis are still incompletely characterized. Bone marrow [CD34⁺, CD38^{neg}, Lin^{neg}] cells contain a HSC population, while CD38⁺ and Lin⁺ fractions contain committed progenitors and mature cells. Within the [CD34⁺, CD38^{neg}, Lin^{neg}] population, HSCs are enriched in the Thy-1⁺ fraction. The origin of Thy-1^{neg} elements within [CD34⁺, CD38^{neg}, Lin^{neg}] cells is uncertain, as is their capacity to self-renew.

the generation of intermediate oligo-lineage precursors (**Figure 1a**). The profile of surface markers used to differentiate and purify mouse HSCs has evolved over time, varies among authors, and is undergoing continuous refinement (10). A detailed description of the mouse HSC surface marker profile is beyond the scope of this article but there are several comprehensive reviews of this subject (6, 11).

Initially developed in mice, the stem cell model of hematopoiesis has been adapted to humans, although the impossibility to perform *in vivo* competitive repopulation assays has limited a detailed functional dissection of all differentiative stages. Most studies on human HSCs have been performed using surrogate *in vitro* and *in vivo* studies, such as clonogenic assays on stromal feeder




Figure 1 (Continued)

(c) In several forms of human acute myeloid leukemia (AML), such as the M1, M4 and M5 variants, the [CD34⁺, CD38^{neg}, Lin^{neg}] fraction contains a leukemic stem cell (LSC) population, but it is not clear whether LSCs are Thy-1⁺ or Thy-1^{neg}. Contrary to normal HSCs, human AML colony-forming unit (CFU) blast cells are usually Thy-1^{neg}. (d) Studies on AML1-ETO⁺ AML provide insights into leukemic hematopoiesis. In patients undergoing complete remission, Thy-1⁺ HSCs harbor the AML1-ETO chromosomal translocation but undergo normal differentiation, indicating that AML1-ETO is not sufficient to induce leukemia. The bone marrow of the same patients at diagnosis is infiltrated by AML1-ETO⁺ CFU-blasts that are [CD34⁺, CD38^{neg}, Lin^{neg}] and Thy-1^{neg}. This observation suggests that full leukemic transformation is likely caused by a second mutation either targeting more differentiated Thy-1^{neg} precursors, or causing loss of Thy-1 expression in LSCs.

layers and repopulation experiments in immunodeficient (SCID, NOD-SCID) mice (6, 12). **Figure 1b** illustrates a simplified, tentative consensus model of human myeloid hematopoiesis. It is widely accepted that the [CD34⁺, CD38^{neg}, Lin^{neg}] fraction of human

bone marrow cells contains a HSC population, whereas the CD38⁺ and Lin⁺ fractions contain more committed progenitors and mature elements (6, 13–15). Within the [CD34⁺, CD38^{neg}, Lin^{neg}] population, HSCs appear particularly enriched in the Thy-1⁺

SCID mouse: severe combined immunodeficiency mouse; an immunodeficient mouse strain, characterized by lack of B and T lymphocytes

NOD-SCID mouse: nonobese diabetic SCID mouse; an immunodeficient mouse strain characterized by lack of B, T, and NK lymphocytes

Hematopoietic stem cell (HSC) mobilization: a procedure by which HSCs are induced to exit the bone marrow and enter blood circulation, usually by cytokine administration

fraction (6, 16), and it is known that transplantation of purified [Thy-1⁺, CD34⁺, Lin^{neg}] cells from peripheral blood after HSC “mobilization” is capable of long-term reconstitution of hematopoiesis in human patients (17). The nature of [Thy-1^{neg}, CD34⁺, CD38^{neg}, Lin^{neg}] cells remains uncertain. It is not clear whether they derive from [Thy-1⁺, CD34⁺, CD38^{neg}, Lin^{neg}] cells, nor if they can self-renew *in vivo*.

Cancer Stem Cells in Human Myeloid Leukemias

In the 1970s, based on the available knowledge of normal hematopoiesis, Fialkow and colleagues began to address the possible relationship between HSCs and human leukemias. They showed that diseases as diverse as chronic myeloid leukemia (CML), acute myeloid leukemia (AML), essential thrombocythemia, and polycythemia vera were all characterized by the expansion of a monoclonal population of cells that contained multiple lineages of differentiated mature blood elements (18–21). In the early 1990s, Dick and colleagues started a series of pioneering investigations to understand whether the functional hierarchy observed in normal hematopoiesis was conserved in blood tumors. These studies showed that in several forms of human AML (M1, M4, M5), the cells with the capacity to engraft in NOD-SCID immunodeficient mice are restricted to a minority subpopulation of the leukemic clone, defined as [CD34⁺, CD38^{neg}] and thus sharing a surface marker profile with normal immature multipotent progenitors (22, 23). Most interestingly, analysis of leukemic cell populations engrafted in NOD-SCID mice revealed reconstitution of the phenotypic heterogeneity observed in the original donor. This indicates that [CD34⁺, CD38^{neg}] leukemic stem cells (LSCs) retained differentiative capacity, giving rise to CD38⁺ and Lin⁺ populations. Taken together, these observations provided the first experimental proof that in a human neoplastic disease (i.e., AML), cancer cell pop-

ulations are organized according to a functional hierarchy mirroring a stem cell system (Figure 1c).

Disease Heterogeneity in Cancer Stem Cell Populations

In addition to providing proof of principle of the CSC model, the study of human AML also shed the first light on its potential complexities. A first observation is that not all forms of AML follow the same paradigm. In patients affected by acute promyelocytic leukemia (APL or AML-M3), the t(15:17) PML/RAR α chromosomal translocation is usually detected in [CD34⁺, CD38⁺] cells, but not in [CD34⁺, CD38^{neg}] cells (24). The APL case is somewhat difficult to judge because of its apparent lack of engraftment ability in NOD-SCID mice and the resulting impossibility of transplantation studies (22). Several explanations could account for this discrepancy with other AML forms (e.g., CD38 expression might be induced in LSCs by tumor transformation). However, the absence of t(15:17) in HSCs of APL patients suggests that, in selected cases, AML growth might be sustained by a cell population that does not originate from HSCs.

“Stem Cell Evolution” During Disease Progression

Another layer of complexity is added by investigations of the “evolution” of LSCs during the natural history of leukemic diseases. A first example is provided by studies performed on AML patients whose disease harbored the t(8:21) AML1-ETO chromosomal translocation and who achieved long-lasting clinical remission (25). In these patients, analysis of bone marrow at remission revealed persistence of AML1-ETO fusion transcripts in [Thy-1⁺, CD34⁺, CD38^{neg}, Lin^{neg}] cells, suggesting that the AML1-ETO chromosomal translocation is an early genetic event targeting the HSC compartment. In addition, *in vitro* functional studies performed on the

same cells showed that AML1-ETO⁺ progenitors were able to regularly differentiate into mature cells of different lineages (**Figure 1d**). Moreover, AML1-ETO fusion transcripts were detectable in multiple lineages of purified mature cells directly obtained from the same patients. Therefore, although the AML1-ETO translocation is probably an early, necessary event in the natural history of this disease, by itself it is not sufficient to cause leukemic transformation. Additional mutations are probably required to induce a fully transformed phenotype, and most likely arise, as a final transforming event, in a subsequent stage of hematopoietic differentiation. Supporting this hypothesis is the observation that, whereas clonogenic assays performed on the bone marrow of AML1-ETO⁺ patients in complete remission reveal a normal repertoire of colony-forming units (CFUs) in the [Thy-1⁺, CD34⁺, CD38^{neg}, Lin^{neg}] population, those performed on AML1-ETO⁺ patients at diagnosis reveal only the presence of “blast”-forming units (CFU-blast) restricted to the [Thy-1^{neg}, CD34⁺, CD38^{neg}, Lin^{neg}] population. This finding, common to most forms of human AML (26), might suggest that the fully transformed LSC, although derived from a mutated preleukemic Thy-1⁺ HSC, could actually emerge as the consequence of secondary mutations in a Thy-1^{neg} more differentiated progenitor (27) (**Figure 1d**).

An intriguingly similar scenario can be envisaged in CML when comparing the initial chronic phase (Cp) of the disease with its terminal stage, blast crisis (Bc). During Cp-CML, the leukemic clone undergoes multipotent differentiation, and the t(9;22) BCR-ABL chromosomal translocation, which serves as a diagnostic marker of the disease, is detectable in most circulating mature lineages, especially myeloid cells and B lymphocytes (28, 29). During Bc-CML, as the disease becomes clinically more aggressive, large numbers of undifferentiated BCR-ABL⁺ blast cells accumulate in the blood and bone marrow. It is not clear which molecular event under-

lies progression from Cp-CML to Bc-CML. However, it has recently been demonstrated that Bc-CML is characterized by an expansion of oligo-lineage precursors, especially the granulocyte-macrophage progenitor (GMP) (30). Most interestingly, when compared to their normal counterparts, GMPs from Bc-CML show aberrant de novo acquisition of in vitro self-renewal properties, associated with increased nuclear expression of β -catenin (30). Thus, during progression from Cp-CML to Bc-CML, a subpopulation of differentiated leukemic progenitors (i.e., GMPs) appears to acquire stem cell characteristics. This observation suggests that, in CML, disease progression is probably accompanied by both an expansion and a diversification of the leukemogenic stem cell pool owing to the generation of a second class of LSCs (31).

Overall, the picture emerging from studies on human leukemias confirms the CSC working model but also highlights the variety of its potential manifestations. The functional hierarchy of tumor cell populations can vary among disease subtypes and can evolve during their natural history, as a cause or as a result of their progression. In many cases, a detailed understanding of the true hierarchy of human leukemic cell populations is still hampered by the many uncertainties regarding the hierarchy of the early stages of normal human hematopoiesis (11). However, the continuous refinement of the surface marker profiles used to distinguish HSCs from MPPs, such as that recently achieved in the mouse using the differential expression of signaling lymphocytic activation molecules (SLAMs) (10), could shed new light on this issue in the near future.

The Molecular Machinery of Self-Renewal in HSCs and LSCs: Similarities and Differences

At a molecular level, applying concepts and tools first developed for the study of normal HSCs to the study of leukemias is leading to new and provocative insights in basic cancer biology. One interesting example is related to

Colony-forming unit (CFU)-blast: a cell that gives rise to aberrant blast colonies when tested in a clonogenic assay for hematopoietic progenitors

GMP (granulocyte-macrophage progenitor): a committed hematopoietic precursor whose progeny is restricted to the monocytic and granulocytic lineages. GMPs derive from CMPs

Bmi1 (B lymphoma Mo-MLV insertion region 1): oncogene identified as a DNA insertion site of the Moloney murine leukemia virus (Mo-MLV)

the study of the Bmi1 oncogene, a member of the Polycomb group ring finger (PCGF) gene family, shown to be expressed at high levels in HSCs and progressively downregulated during hematopoietic differentiation (32). Bmi1 knockout (Bmi1^{-/-}) mice are characterized by progressive bone marrow aplasia and die less than two months after birth (33). In this system, progressive bone marrow aplasia is not associated with differentiation problems but with progressive numerical reduction of bone marrow HSCs. Most interestingly, when transplanted in syngeneic wild-type mice, both fetal liver and bone marrow HSCs obtained from Bmi1^{-/-} mice sustain hematopoiesis in a short-term, transient fashion, thus indicating that Bmi1^{-/-} HSCs lack self-renewal capacity because of an intrinsic cell defect and behave functionally as MPPs (33).

When assessed in a murine leukemia model, the role of Bmi1 in self-renewal has proven to have key implications for tumor biology. In mice, infection of fetal liver HSCs with a retroviral construct simultaneously encoding Hoxa9 and Meis1 oncogenes is sufficient to induce tumor transformation and generation of a leukemic disease resembling human AML (34). When this experiment is performed using Bmi1^{-/-} HSCs, no substantial difference is initially observed: the virus construct induces tumor transformation and generation of a Bmi1^{-/-} AML that rapidly kills the recipient mice (35). However, when the long-term self-renewal of leukemic cells is evaluated by serial transplantation in secondary recipients, only Bmi1^{+/+} AML cells are able to transfer the disease. Bmi1^{-/-} AML cells cannot, thus indicating that lack of Bmi1, while not interfering with the tumor transformation process per se, limits the expansion potential of the leukemic clone (35).

A second molecule that is likely to play a key role in the molecular machinery of both HSC and LSC self-renewal is the protein phosphatase and tensin homologue (PTEN), a known tumor suppressor (36). In mice, conditional deletion of the Pten gene causes pro-

gressive depletion of the HSC compartment without substantially interfering with its differentiation capacity, a scenario reminiscent of Bmi1^{-/-} mice (37, 38). However, in contrast to the case of Bmi1, inactivation of Pten is also associated with spontaneous development of acute leukemias, lethal in both primary hosts and secondary transplant recipients (37, 38).

Taken together, these studies show that genes required for self-renewal of normal HSCs can play opposite roles in the development of leukemia. In some cases they are necessary for long-term expansion of the transformed clone (Bmi1), but in others they act as tumor suppressors and prevent leukemic transformation (Pten). Most interestingly, these studies show that two of the common hallmarks of cancer—the endowment of long-term proliferative potential (historically sometimes referred to as “immortality”) and the propensity toward uncontrolled, aberrant growth (“transformation”)—are two independent properties that can be dissociated, a concept also indirectly demonstrated by in vitro studies on primary cultures of human tumors (39).

CANCER STEM CELLS IN HUMAN SOLID TUMORS

Several tissues share with blood the necessity to undergo a continuous or cyclical renovation. Among them are the skin and all major epithelia of the gastrointestinal (mouth, pharynx, esophagus, stomach, gut), respiratory (larynx, trachea, bronchi, lungs), and reproductive and genitourinary (breast, ovary, vagina, uterus, bladder, prostate) systems. Most of them are known or presumed to be organized according to a hierarchical system based on a dedicated adult stem cell population (40–52). Moreover, many human tissues traditionally considered stable, undergoing only minimal or slow turnover throughout adult life, such as the brain and muscle, are today known to contain specific stem cell populations, which can be mobilized and

activated in specific situations (53, 54). In addition, a stem-like functional hierarchy, although described using a different terminology, is known to exist in the mature lymphoid system: T and B cell populations can be subdivided into long-term (“memory”) and short-term (“effector”) functional compartments, characterized by specific surface marker expression profiles (55).

On the basis of these observations, our laboratory undertook a new study to test whether a CSC model could be applied to the description of solid tumors, focusing on human breast cancer as a model system (56). The results showed that, in most human breast cancers, only a minority subpopulation of the tumor clone, defined as [CD44⁺, CD24^{-/low}] and representing 11%–35% of total cancer cells, is endowed with the capacity to sustain tumor growth when xenografted in NOD-SCID mice. Most importantly, tumors grown from [CD44⁺, CD24^{-/low}] cells were shown to contain mixed populations of epithelial tumor cells, recreating the phenotypic heterogeneity of the parent tumors. For the first time, this study demonstrated the existence of a functional hierarchy reminiscent of stem cell systems in a solid human epithelial tumor. The limited knowledge of the biology of normal human mammary stem cells and their differentiation programs (57) currently limits our understanding of which lineage, progenitor cell, or differentiation stage of the mammary epithelial tissue is related to [CD44⁺, CD24^{-/low}] human breast CSC. However, development of *in vitro* culture systems for the functional study of both normal (58) and cancer (59) human mammary epithelial cells could soon provide the experimental means for a phenotypic dissection of breast epithelial differentiation processes.

The CSC working model has also been successfully applied to brain tumors (60–63). Studies performed on glioblastoma multiforme and medulloblastoma have shown that tumorigenic cells are restricted to the CD133⁺ subpopulation, which usually represents 5%–30% of total tumor cells. As ex-

pected from the CSC model, tumors resulting from orthotopic, intracerebral injection of CD133⁺ cells reproduced the phenotypic diversity and differentiation pattern of the parent tumors (62). In the study of brain tumors, the availability of a well-characterized cell culture system for normal neural stem cells (the “neurosphere” assay) provides a robust tool for the *in vitro* study of their candidate pathological counterparts. Based on this approach, Galli et al. succeeded in the isolation and serial propagation from human glioblastoma multiforme of “cancer neurospheres,” which are highly enriched in long-term self-renewing, multi-lineage-differentiating, and tumor-initiating cells (60). Although the relationship between CD133⁺ cells and cancer neurospheres still remains to be fully explored, it is safe to assume that *ex vivo* purification of brain tumor CSCs based on CD133 coupled with *in vitro* functional studies using neurosphere assays will provide one of the most effective probes for the study of solid-tumor CSCs in the near future.

In addition to breast and brain tumors, interesting results are being accumulated on prostate cancer, where progenitor/tumorigenic subpopulations appear characterized by the expression of CD44 (64, 65). In melanoma, *in vitro* culture of primary tumor cells with media formulations used for embryonic stem cells allows the isolation and serial propagation of “melanoma spheroids,” reminiscent of cancer neurospheres and capable of long-term self-renewal, multi-lineage differentiation and *in vivo* tumorigenicity (66). Several mouse models for the study of CSC biology in epithelial solid tumors are also currently being investigated, including models for lung (48) and prostate (67) cancer.

The CSC model is also being extended to lymphoid tumors, such as multiple myeloma, where it is becoming apparent that a minority subpopulation of CD138^{neg} cells is preferentially endowed with *in vitro* clonogenic and *in vivo* engraftment capacity (68). Studies on multiple myeloma are particularly fascinating because they suggest that the

candidate CSC for plasma cell diseases might be a B cell population in an earlier stage of differentiation, which might reside in a different anatomical compartment (i.e., lymph nodes) than the bulk of the tumor (i.e., mature CD138⁺ plasma cells in the bone marrow).

IMPLICATIONS OF THE CSC MODEL FOR THE BIOLOGICAL INTERPRETATION OF CANCER

Intratumor Heterogeneity

One conceptual implication of the CSC model is related to the biological interpretation of intratumor heterogeneity, i.e., heterogeneity among cancer cells within the same tumor lesion. Intratumor heterogeneity is a recurrent observation in human tumors, both primary and metastatic (69), and is usually considered the result of variations in the tumor microenvironment and the coexistence of multiple independent cellular subclones originated by progressive accumulation of divergent genetic aberrations within the same cancer cell population (70, 71). The CSC model introduces an additional explanation, suggesting that intratumor heterogeneity can also result from functional diversity among cells in different states of differentiation.

This account has major theoretical and methodological consequences for cancer research because it suggests that the biological significance of observations collected on whole tumor tissues should be judged according to their relation to the CSC subset. For instance, biochemical pathways that are active in the majority of tumor cells might be of little functional relevance for the biology of CSC, whereas biochemical pathways active only on a small minority of cancer cells might play key roles in CSC biology and thus in the overall long-term behavior of a tumor.

Based on this perspective, the CSC model might help shed light on several unexpected observations in cancer biology. An interesting example is the heterogeneous and frequently nonconstitutive pattern of intratumor expres-

sion of the human telomerase reverse transcriptase (hTERT) (39, 72). According to current models, based mainly on the study of in vitro cultured cell lines, one of the hallmarks of cancer is the acquisition by cancer cells of a limitless replicative potential, or immortality (73). To become immortal, cancer cells must be endowed with a system for the maintenance of telomere length, in most cases based on telomerase, a ribonucleoproteic enzymatic complex that counteracts the progressive shortening of telomere repeat sequences during in vitro proliferation of normal somatic cells (74). Because telomerase expression is usually not detected in most normal tissues, it is postulated that its expression in cancer cells is the result of tumor-specific genetic and/or epigenetic events that result in ectopic, constitutive hTERT activation (75). Despite extensive investigation, however, the molecular mechanisms underlying telomerase upregulation remain largely undefined (75), and contrary to predictions, the in vivo expression of hTERT has repeatedly proven to be extremely heterogeneous among cancer cells, especially in vivo (39, 72). These observations can be explained in the context of the CSC model if we assume that telomerase expression is restricted to the self-renewing, long-lived, tumorigenic subpopulation of the tumor clone (i.e., CSC). According to this hypothesis, telomerase expression is inherited by CSCs as part of the normal self-renewal machinery of adult stem cells of the corresponding healthy parent tissues and is progressively lost during differentiation processes, both in normal and in tumor tissues (76). Indeed, it has recently been shown that one of the genes involved in the control of stem cell self-renewal, *Bmi1*, is also able to upregulate hTERT expression in epithelial cells (77). Moreover, it is now well known that telomerase is expressed in normal adult stem cells of healthy tissues, although at levels that are usually below the threshold for routine experimental detection (72, 78). The scenario is probably similar for other proteins that are known to display similar expression patterns

and to play key roles in tumor biology, such as survivin (69, 79).

CSCs and the Study of Tumor Metastasis

The CSC model can also shed new light on the biology of metastases and explain why, despite extensive intratumor heterogeneity (69, 71), comparison of paired samples of primary tumors and autologous lymph node and/or distant-site metastases usually reveals striking similarities over a wide range of parameters, including tissue morphology (69, 71), repertoire of somatic genetic mutations (80–82), expression of tumor-suppressor and immunomodulatory proteins (83), expression of epigenetically controlled genes (84), and overall transcriptional profile as defined by gene-expression arrays (85–87). These observations are in contrast to predictions from traditional cancer models, where metastases are considered to originate from monoclonal expansions of very specific, individual tumor subclones endowed with specific genotypic and phenotypic features, and therefore are postulated to be substantially different from primary tumors (**Figure 2a**). However, if we take into account the CSC model and we assume that, in each individual tumor, the differentiation pattern is controlled by its specific repertoire of genetic mutations, then we can predict that, if two lesions share identical genetic backgrounds and similar genetic abnormalities, they will also undergo similar differentiation programs and display similar patterns of intratumor heterogeneity in the expression of differentiation antigens (**Figure 2b**) (69, 87).

IMPLICATIONS OF THE CSC MODEL FOR DESIGN AND EVALUATION OF ANTITUMOR TREATMENTS

Therapeutic Tumor Targeting

The observation that cancer growth can be sustained by a minority subpopulation of tu-

mor cells with unique functional properties (i.e., CSCs) could also assist in the design of new and more effective antitumor treatments. According to the CSC model, therapeutic approaches that do not eradicate the CSC compartment are likely to achieve little success; they might kill the majority of tumor cells and induce temporary regression of gross tumor lesions but fail to prevent disease relapse and metastatic dissemination (**Figure 3a**) (2). In support of this hypothesis is the finding that, in the hematopoietic system, both normal stem cells and CSCs (i.e., HSCs and LSCs from AML patients) mainly appear to be in a quiescent, nondividing, G_0 state, and therefore inherently resistant to the toxic effect of traditional chemotherapeutic regimens (8, 88, 89). Based on this concept, traditional treatments might be recalibrated and investigational therapies developed, focusing on their ability to target the CSC subpopulation and its specific biochemical pathways (89, 90). Similarly, the CSC model might explain why several experimental therapeutic approaches have shown poor clinical results despite extensive preclinical validation *in vitro*, and it might provide critical information for the redesign and upgrading of such approaches. A classic example of this situation could be represented by antitumor T cell immunotherapies, where target antigen selection might be re-evaluated on the basis of expression on CSCs (91, 92).

Evaluation of Treatment Efficacy

A second therapeutic implication of the CSC model is related to clinical methodologies for evaluating treatment efficacy. Traditionally, antitumor treatments are screened based on their capacity to induce a clinical response (i.e., a dramatic regression, either complete or partial, of the tumor lesion). This approach, however, tends to select for treatments that are active on the bulk of tumor cell populations but not necessarily on CSCs. From a purely theoretical point of view, antitumor treatments that selectively target the CSC subset might actually be unable to




Figure 2

Impact of the cancer stem cell (CSC) model on the origin and biology of metastases. (a) According to standard cancer models, tumors are composed of heterogeneous mixtures of independent subclones, originated by divergent genetic mutations; different subclones are endowed with different functional properties, and only selected clones (*dark grey cells*) can migrate and form metastases. The metastasis is predicted to be a homogeneous monoclonal expansion of an individual subclone, which in turn can accumulate further mutations (*striped and variously patterned cells*) and diverge even further from the primary tumor. Overall, the model predicts that primary tumors and corresponding metastases are substantially different. (b) The CSC model assumes that intratumor heterogeneity is mainly caused by cell differentiation, and that only CSCs (*dark grey cells*) can migrate and form overt metastases, while differentiated cells (*light grey cells*) undergo apoptosis. In the CSC model, metastatic cancer tissues undergo differentiation programs that closely resemble those observed in the corresponding primary tissues. Recent experimental evidence based on gene-expression microarrays tends to support the CSC model for human epithelial tumors, such as breast and colon cancer. The two hypotheses are not mutually exclusive, and elements of both are probably true.




Figure 3

Impact of the cancer stem cell (CSC) model on the design and evaluation of antitumor treatments. (a) Antitumor treatments designed and selected for broad cytotoxic activity may kill the majority of cancer cells within a specific tumor tissue and induce dramatic, even complete regression of large tumor masses; however, if any of the CSCs are spared, tumor tissues can be regenerated and cause disease relapse. (b) In contrast, antitumor treatments specifically designed to target CSCs, although theoretically unable to cause rapid shrinkage of tumor lesions, might nonetheless achieve long-term disease eradication by exhausting self-renewal and growth potential of cancer tissues.

induce rapid shrinkage of tumor masses but might eliminate their capacity for long-term growth and therefore cause their arrest or slowly reduce their size (Figure 3b). It is

therefore likely that, alongside new treatment strategies, new approaches for the preclinical evaluation of their efficacy will need to be devised.

DISCLOSURE STATEMENT

MFC owns stock in Oncomed, a cancer stem cell company.

ACKNOWLEDGMENTS

We thank Drs. Michael W. Becker (University of Rochester, Rochester, NY), Laurie Ailles, Joanne Attema and Ravi Majeti (Stanford University, Stanford, CA) for helpful suggestions and

critical discussions. P.D. was supported by the Leonino Fontana e Maria Lionello fellowship from Fondazione Italiana per la Ricerca sul Cancro (FIRC) and subsequently by a fellowship from the California Institute of Regenerative Medicine (CIRM) training program. This work was supported by NIH grant CA104987-02.

LITERATURE CITED

- Clarke MF, Fuller M. 2006. Stem cells and cancer: two faces of Eve. *Cell* 124:1111–15
- Reya T, Morrison SJ, Clarke MF, Weissman IL. 2001. Stem cells, cancer, and cancer stem cells. *Nature* 414:105–11
- Fearon ER, Vogelstein B. 1990. A genetic model for colorectal tumorigenesis. *Cell* 61:759–67
- Clarke MF. 2005. A self-renewal assay for cancer stem cells. *Cancer Chemother. Pharmacol.* 56(Suppl. 1):64–68
- Till JE, McCulloch EA. 1961. A direct measurement of the radiation sensitivity of normal mouse bone marrow cells. *Radiat. Res.* 14:213–22
- Kondo M, Wagers AJ, Manz MG, et al. 2003. Biology of hematopoietic stem cells and progenitors: implications for clinical application. *Annu. Rev. Immunol.* 21:759–806
- Spangrude GJ, Heimfeld S, Weissman IL. 1988. Purification and characterization of mouse hematopoietic stem cells. *Science* 241:58–62
- Morrison SJ, Weissman IL. 1994. The long-term repopulating subset of hematopoietic stem cells is deterministic and isolatable by phenotype. *Immunity* 1:661–73
- Morrison SJ, Wandycz AM, Hemmati HD, et al. 1997. Identification of a lineage of multipotent hematopoietic progenitors. *Development* 124:1929–39
- Kiel MJ, Yilmaz OH, Iwashita T, et al. 2005. SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells. *Cell* 121:1109–21
- Passegue E, Jamieson CH, Ailles LE, Weissman IL. 2003. Normal and leukemic hematopoiesis: Are leukemias a stem cell disorder or a reacquisition of stem cell characteristics? *Proc. Natl. Acad. Sci. USA* 100(Suppl. 1):11842–49
- Dick JE, Bhatia M, Gan O, et al. 1997. Assay of human stem cells by repopulation of NOD/SCID mice. *Stem Cells* 15(Suppl. 1):199–203; discussion 204–7
- Bhatia M, Wang JC, Kapp U, et al. 1997. Purification of primitive human hematopoietic cells capable of repopulating immune-deficient mice. *Proc. Natl. Acad. Sci. USA* 94:5320–25
- Manz MG, Miyamoto T, Akashi K, Weissman IL. 2002. Prospective isolation of human clonogenic common myeloid progenitors. *Proc. Natl. Acad. Sci. USA* 99:11872–77
- Terstappen LW, Huang S, Safford M, et al. 1991. Sequential generations of hematopoietic colonies derived from single nonlineage-committed CD34+CD38– progenitor cells. *Blood* 77:1218–27
- Baum CM, Weissman IL, Tsukamoto AS, et al. 1992. Isolation of a candidate human hematopoietic stem-cell population. *Proc. Natl. Acad. Sci. USA* 89:2804–8
- Shizuru JA, Negrin RS, Weissman IL. 2005. Hematopoietic stem and progenitor cells: clinical and preclinical regeneration of the hematolymphoid system. *Annu. Rev. Med.* 56:509–38
- Fialkow PJ, Faguet GB, Jacobson RJ, et al. 1981. Evidence that essential thrombocythemia is a clonal disorder with origin in a multipotent stem cell. *Blood* 58:916–19
- Fialkow PJ, Jacobson RJ, Papayannopoulou T. 1977. Chronic myelocytic leukemia: clonal origin in a stem cell common to the granulocyte, erythrocyte, platelet and monocyte/macrophage. *Am. J. Med.* 63:125–30

19. Among the first studies to provide evidence of the stem cell origin of a human hematologic malignancy.

20. Adamson JW, Fialkow PJ, Murphy S, et al. 1976. Polycythemia vera: stem-cell and probable clonal origin of the disease. *N. Engl. J. Med.* 295:913–16
21. Fialkow PJ, Singer JW, Adamson JW, et al. 1981. Acute nonlymphocytic leukemia: heterogeneity of stem cell origin. *Blood* 57:1068–73
22. **Bonnet D, Dick JE. 1997. Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nat. Med.* 3:730–37**
23. Lapidot T, Sirard C, Vormoor J, et al. 1994. A cell initiating human acute myeloid leukaemia after transplantation into SCID mice. *Nature* 367:645–48
24. Turhan AG, Lemoine FM, Debert C, et al. 1995. Highly purified primitive hematopoietic stem cells are PML-RARA negative and generate nonclonal progenitors in acute promyelocytic leukemia. *Blood* 85:2154–61
25. **Miyamoto T, Weissman IL, Akashi K. 2000. AML1/ETO-expressing nonleukemic stem cells in acute myelogenous leukemia with 8;21 chromosomal translocation. *Proc. Natl. Acad. Sci. USA* 97:7521–26**
26. Blair A, Hogge DE, Ailles LE, et al. 1997. Lack of expression of Thy-1 (CD90) on acute myeloid leukemia cells with long-term proliferative ability in vitro and in vivo. *Blood* 89:3104–12
27. Weissman IL. 2005. Stem cell research: paths to cancer therapies and regenerative medicine. *JAMA* 294:1359–66
28. Takahashi N, Miura I, Saitoh K, Miura AB. 1998. Lineage involvement of stem cells bearing the Philadelphia chromosome in chronic myeloid leukemia in the chronic phase as shown by a combination of fluorescence-activated cell sorting and fluorescence in situ hybridization. *Blood* 92:4758–63
29. Primo D, Sanchez ML, Espinosa AB, et al. 2006. Lineage involvement in chronic myeloid leukaemia: comparison between MBCR/ABL and mBCR/ABL cases. *Br. J. Haematol.* 132:736–9
30. Jamieson CH, Ailles LE, Dylla SJ, et al. 2004. Granulocyte-macrophage progenitors as candidate leukemic stem cells in blast-crisis CML. *N. Engl. J. Med.* 351:657–67
31. Clarke MF. 2004. Chronic myelogenous leukemia—identifying the hydra’s heads. *N. Engl. J. Med.* 351:634–36
32. Lessard J, Baban S, Sauvageau G. 1998. Stage-specific expression of polycomb group genes in human bone marrow cells. *Blood* 91:1216–24
33. **Park IK, Qian D, Kiel M, et al. 2003. Bmi-1 is required for maintenance of adult self-renewing hematopoietic stem cells. *Nature* 423:302–5**
34. Kroon E, Kros J, Thorsteinsdottir U, et al. 1998. Hoxa9 transforms primary bone marrow cells through specific collaboration with Meis1a but not Pbx1b. *EMBO J.* 17:3714–25
35. **Lessard J, Sauvageau G. 2003. Bmi-1 determines the proliferative capacity of normal and leukaemic stem cells. *Nature* 423:255–60**
36. Di Cristofano A, Pandolfi PP. 2000. The multiple roles of PTEN in tumor suppression. *Cell* 100:387–90
37. Yilmaz OH, Valdez R, Theisen BK, et al. 2006. Pten dependence distinguishes hematopoietic stem cells from leukaemia-initiating cells. *Nature* 441:475–82
38. Zhang J, Grindley JC, Yin T, et al. 2006. PTEN maintains hematopoietic stem cells and acts in lineage choice and leukaemia prevention. *Nature* 441:518–22
39. **Dalerba P, Guiducci C, Poliani PL, et al. 2005. Reconstitution of human telomerase reverse transcriptase expression rescues colorectal carcinoma cells from in vitro senescence: evidence against immortality as a constitutive trait of tumor cells. *Cancer Res.* 65:2321–29**

22. Seminal study describing the first formal isolation of a CSC population in a human neoplastic disease (AML).

25. First study showing that CSC populations can change during disease progression in a human hematologic malignancy.

33. First identification of a gene required for the maintenance of adult stem cell self-renewal.

35. First dissection of self-renewal as an independent feature required for long-term expansion of cancer cells.

39. First report providing functional evidence of the nonconstitutive nature of telomerase activation in a human cancer.

40. Shackleton M, Vaillant F, Simpson KJ, et al. 2006. Generation of a functional mammary gland from a single stem cell. *Nature* 439:84–88
41. Stingl J, Eirew P, Ricketson I, et al. 2006. Purification and unique properties of mammary epithelial stem cells. *Nature* 439:993–97
42. Brittan M, Wright NA. 2004. Stem cell in gastrointestinal structure and neoplastic development. *Gut* 53:899–910
43. Bach SP, Renahan AG, Potten CS. 2000. Stem cells: the intestinal stem cell as a paradigm. *Carcinogenesis* 21:469–76
44. Seery JP. 2002. Stem cells of the esophageal epithelium. *J. Cell Sci.* 115:1783–89
45. Watt FM. 2001. Stem cell fate and patterning in mammalian epidermis. *Curr. Opin. Genet. Dev.* 11:410–17
46. Sancho E, Battle E, Clevers H. 2003. Live and let die in the intestinal epithelium. *Curr. Opin. Cell Biol.* 15:763–70
47. Alonso L, Fuchs E. 2003. Stem cells of the skin epithelium. *Proc. Natl. Acad. Sci. USA* 100(Suppl. 1):11830–35
48. Kim CF, Jackson EL, Woolfenden AE, et al. 2005. Identification of bronchioalveolar stem cells in normal lung and lung cancer. *Cell* 121:823–35
49. Otto WR. 2002. Lung epithelial stem cells. *J. Pathol.* 197:527–35
50. Gargett CE. 2004. Stem cells in gynaecology. *Aust. NZ J. Obstet. Gynaecol.* 44:380–86
51. Potten CS. 1998. Stem cells in gastrointestinal epithelium: numbers, characteristics and death. *Philos. Trans. R. Soc. London Ser. B* 353:821–30
52. Schalken JA, van Leenders G. 2003. Cellular and molecular biology of the prostate: stem cell biology. *Urology* 62:11–20
53. Ming GL, Song H. 2005. Adult neurogenesis in the mammalian central nervous system. *Annu. Rev. Neurosci.* 28:223–50
54. Collins CA, Olsen I, Zammit PS, et al. 2005. Stem cell function, self-renewal, and behavioral heterogeneity of cells from the adult muscle satellite cell niche. *Cell* 122:289–301
55. Luckey CJ, Bhattacharya D, Goldrath AW, et al. 2006. Memory T and memory B cells share a transcriptional program of self-renewal with long-term hematopoietic stem cells. *Proc. Natl. Acad. Sci. USA* 103:3304–9
56. Al-Hajj M, Wicha MS, Benito-Hernandez A, et al. 2003. **Prospective identification of tumorigenic breast cancer cells.** *Proc. Natl. Acad. Sci. USA* 100:3983–88
57. Stingl J, Raouf A, Emerman JT, Eaves CJ. 2005. Epithelial progenitors in the normal human mammary gland. *J. Mammary Gland Biol. Neoplasia* 10:49–59
58. Dontu G, Abdallah WM, Foley JM, et al. 2003. In vitro propagation and transcriptional profiling of human mammary stem/progenitor cells. *Genes Dev.* 17:1253–70
59. Ponti D, Costa A, Zaffaroni N, et al. 2005. Isolation and in vitro propagation of tumorigenic breast cancer cells with stem/progenitor cell properties. *Cancer Res.* 65:5506–11
60. Galli R, Binda E, Orfanelli U, et al. 2004. Isolation and characterization of tumorigenic, stem-like neural precursors from human glioblastoma. *Cancer Res.* 64:7011–21
61. Singh SK, Clarke ID, Terasaki M, et al. 2003. Identification of a cancer stem cell in human brain tumors. *Cancer Res.* 63:5821–28
62. Singh SK, Hawkins C, Clarke ID, et al. 2004. Identification of human brain tumor initiating cells. *Nature* 432:396–401
63. Vescovi AL, Galli R, Reynolds BA. 2006. Brain tumor stem cells. *Nat. Rev. Cancer* 6:425–36
64. Collins AT, Berry PA, Hyde C, et al. 2005. Prospective identification of tumorigenic prostate cancer stem cells. *Cancer Res.* 65:10946–51

56. First report describing isolation of a CSC population in a human solid tumor.

65. Patrawala L, Calhoun T, Schneider-Broussard R, et al. 2006. Highly purified CD44+ prostate cancer cells from xenograft human tumors are enriched in tumorigenic and metastatic progenitor cells. *Oncogene* 25:1696–708
66. Fang D, Nguyen TK, Leishear K, et al. 2005. A tumorigenic subpopulation with stem cell properties in melanomas. *Cancer Res.* 65:9328–37
67. Xin L, Lawson DA, Witte ON. 2005. The Sca-1 cell surface marker enriches for a prostate-regenerating cell subpopulation that can initiate prostate tumorigenesis. *Proc. Natl. Acad. Sci. USA* 102:6942–47
68. Matsui W, Huff CA, Wang Q, et al. 2004. Characterization of clonogenic multiple myeloma cells. *Blood* 103:2332–36
69. Brabletz T, Jung A, Spaderna S, et al. 2005. Opinion: migrating cancer stem cells—an integrated concept of malignant tumor progression. *Nat. Rev. Cancer* 5:744–49
70. Losi L, Baisse B, Bouzourene H, Benhattar J. 2005. Evolution of intratumoral genetic heterogeneity during colorectal cancer progression. *Carcinogenesis* 26:916–22
71. Brabletz T, Jung A, Reu S, et al. 2001. Variable beta-catenin expression in colorectal cancers indicates tumor progression driven by the tumor environment. *Proc. Natl. Acad. Sci. USA* 98:10356–61
72. Yan P, Benhattar J, Seelentag W, et al. 2004. Immunohistochemical localization of hTERT protein in human tissues. *Histochem. Cell Biol.* 121:391–97
73. Hanahan D, Weinberg RA. 2000. The hallmarks of cancer. *Cell* 100:57–70
74. Shay JW, Wright WE. 2005. Senescence and immortalization: role of telomeres and telomerase. *Carcinogenesis* 26:867–74
75. Kyo S, Inoue M. 2002. Complex regulatory mechanisms of telomerase activity in normal and cancer cells: How can we apply them for cancer therapy? *Oncogene* 21:688–97
76. Morrison SJ, Prowse KR, Ho P, Weissman IL. 1996. Telomerase activity in hematopoietic cells is associated with self-renewal potential. *Immunity* 5:207–16
77. Dimri GP, Martinez JL, Jacobs JLL, et al. 2002. The Bmi-1 oncogene induces telomerase activity and immortalizes human mammary epithelial cells. *Cancer Res.* 62:4736–45
78. Yan P, Saraga EP, Bouzourene H, et al. 2001. Expression of telomerase genes correlates with telomerase activity in human colorectal carcinogenesis. *J. Pathol.* 193:21–26
79. Altieri DC. 2003. Survivin, versatile modulation of cell division and apoptosis in cancer. *Oncogene* 22:8581–89
80. Losi L, Benhattar J, Costa J. 1992. Stability of K-ras mutations throughout the natural history of human colorectal cancer. *Eur. J. Cancer* 28A:1115–20
81. Khan ZA, Jonas SK, Le-Marer N, et al. 2000. P53 mutations in primary and metastatic tumors and circulating tumor cells from colorectal carcinoma patients. *Clin. Cancer Res.* 6:3499–504
82. Zauber P, Sabbath-Solitare M, Marotta SP, Bishop DT. 2003. Molecular changes in the K-ras and APC genes in primary colorectal carcinoma and synchronous metastases compared with the findings in accompanying adenomas. *Mol. Pathol.* 56:137–40
83. Menon AG, Tollenaar RA, van de Velde CJ, et al. 2004. p53 and HLA class-I expression are not down-regulated in colorectal cancer liver metastases. *Clin. Exp. Metastasis* 21:79–85
84. Dalerba P, Ricci A, Russo V, et al. 1998. High homogeneity of MAGE, BAGE, GAGE, tyrosinase and Melan-A/MART-1 gene expression in clusters of multiple simultaneous metastases of human melanoma: implications for protocol design of therapeutic antigen-specific vaccination strategies. *Int. J. Cancer* 77:200–4
85. D'Arrigo A, Belluco C, Ambrosi A, et al. 2005. Metastatic transcriptional pattern revealed by gene expression profiling in primary colorectal carcinoma. *Int. J. Cancer* 115:256–62

86. First description of conserved transcriptional profiles among primary tumors and autologous metastases in breast cancer patients.

86. Weigelt B, Glas AM, Wessels LF, et al. 2003. Gene expression profiles of primary breast tumors maintained in distant metastases. *Proc. Natl. Acad. Sci. USA* 100:15901–5
87. Weigelt B, Peterse JL, van 't Veer LJ. 2005. Breast cancer metastasis: markers and models. *Nat. Rev. Cancer* 5:591–602
88. Guan Y, Gerhard B, Hogge DE. 2003. Detection, isolation, and stimulation of quiescent primitive leukemic progenitor cells from patients with acute myeloid leukemia (AML). *Blood* 101:3142–49
89. Guzman ML, Swiderski CF, Howard DS, et al. 2002. Preferential induction of apoptosis for primary human leukemic stem cells. *Proc. Natl. Acad. Sci. USA* 99:16220–25
90. Al-Hajj M, Becker MW, Wicha M, et al. 2004. Therapeutic implications of cancer stem cells. *Curr. Opin. Genet. Dev.* 14:43–47
91. Dalerba P, Maccalli C, Casati C, et al. 2003. Immunology and immunotherapy of colorectal cancer. *Crit. Rev. Oncol. Hematol.* 46:33–57
92. Parmiani G, Castelli C, Dalerba P, et al. 2002. Cancer immunotherapy with peptide-based vaccines: What have we achieved? Where are we going? *J. Natl. Cancer Inst.* 94:805–18

Michael F. Clarke

A self-renewal assay for cancer stem cells

Published online: 5 November 2005
© Springer-Verlag 2005

Abstract Cancers of epithelial origin are responsible for the majority of cancer-related deaths in the USA. Unfortunately, although chemotherapy and/or radiation therapy can sometimes shrink tumors, metastatic cancers of epithelial origin are essentially incurable. It is clear that new approaches are needed to treat these diseases. Although cancer cell lines provide invaluable information, their biological properties often differ in crucial ways from de novo cancer cells. Our laboratory has developed a novel mouse model that reliably permits individual cancer cells isolated directly from patients' tumors to be assayed. This will allow the characterization of crucial signaling pathways involved in processes such as self-renewal that are critical for tumor formation by the cancer cells within de novo tumors. These tools should lead to new insights into the cellular and molecular mechanisms that drive human breast cancer growth and invasion.

Keywords Stem cells · Cancer · Self-renewal

Introduction

Although cell lines have led to remarkable advances in our understanding of the molecular and biochemical changes in cancer cells, their use in the identification of effective cancer therapies is somewhat limited [5, 15]. Cell lines are imperfect predictors of drug efficacy in de novo tumors. Several factors likely account for this deficiency. Cancer cell lines are selected from a sub-

population of cancer cells that are specifically adapted to growth in tissue culture, and the biological and functional properties of these cell lines can change dramatically [8, 16, 20, 34]. Furthermore, cancer cells from only a minority of breast cancer tumors will establish cell lines, suggesting a selection bias for cancers capable of growing in tissue culture. Additionally, the phenotypic and functional characteristics of these cell lines can change drastically relative to their properties in vivo. For example, the marker expression of both normal hematopoietic and leukemia tissue culture cells can change rapidly in tissue culture and often does not reflect that of the original stem cells from which they were derived. Even when conditions are devised to permit the proliferation of normal stem cells in culture, the conditions often promote self-renewal or differentiation in a way that prevents the stem cells in culture from recapitulating the hierarchy of cell populations that exist in vivo. Taken together, these observations suggest that the biological properties of cancer cell lines can differ markedly from the cancer cells from which they were derived. This likely explains at least in part why the cell lines are often poor predictors of a drug's efficacy in the clinic.

An immunodeficient mouse model for solid tumors

Realizing that a reliable assay was necessary to study the self-renewal of dissociated cancer cells, a substantial and ultimately successful effort by our laboratory led to the development of a xenograft assay that measured not just proliferation but also cancer cell self-renewal [2]. This assay appears to be reliable in permitting dissociated cancer cells from patients' tumors to self-renew. The success of this assay is significantly better than tissue-culture assays and other animal models reported to date. Although breast cancer cells often briefly proliferate in certain tissue-culture systems, cells from most tumors do not self-renew in vitro and eventually stop proliferating [9, 10, 18]. Previously developed mouse models were not

This work was presented at the 20th Bristol-Myers Squibb Nagoya International Cancer Treatment Symposium, "New Concepts of Treatment Strategies for Hormone-Related Cancer", 11–12 March 2005, Nagoya, Japan.

M. F. Clarke
University of Michigan Medical School, 1500 East Medical Center Drive, CCGC 4303, Ann Arbor, MI 48109, USA
E-mail: mclarke@umich.edu
Tel.: +1-734-7648195
Fax: +1-734-6479654

efficient when dissociated cells were injected into mice. This model has recently been extended to other epithelial tumors and we have found that, like breast cancer cells, only a subset of the cancer cells was able to form tumors.

Using this model, we found that only a subset of cancer cells isolated from breast tumors was able to form tumors. Using flow cytometry, we found that the $CD24^{-/low} CD44^{+}$ cancer cells were highly enriched such that cancer cells were able to form tumors in the xenograft model (Fig. 1). On the other hand, in most patients who were examined, other populations of cancer cells were depleted of tumor-forming cells.

Stem cells, cancer and self-renewal

Common cancers arise in tissues that contain a large subpopulation of proliferating cells that are responsible for replenishing the short-lived mature cells. In such organs, cell maturation is arranged in a hierarchy in which a rare population of stem cells gives rise to the mature cells that perpetuate themselves through a process called self-renewal [1, 4, 6, 12, 19, 24, 26, 29, 35]. Owing to their rarity, stem cells must be isolated in order to study their biological, molecular and biochemical properties. Several aspects of stem-cell biology are relevant to cancer. First, both normal stem cells and cancer stem cells undergo self-renewal, and emerging evidence suggests that similar molecular mechanisms regulate self-renewal in normal stem cells and their malignant counterparts. Next, it is likely that mutations that lead to cancer accumulate in normal stem cells. Finally, it

appears that tumors contain a “cancer stem cell” population with indefinite proliferative potential that drives the growth and metastasis of tumors [2].

Genetic regulation of self-renewal in normal stem cells and cancer cells

Maintenance of a tissue or a tumor is determined by a balance of proliferation and cell death [13]. In a normal tissue, stem cell numbers are under tight genetic regulation resulting in a constant number of stem cells in the organ [21, 22, 25]. By contrast, cancer cells have escaped this homeostatic regulation, and the number of cells within a tumor that have the ability to self-renew is constantly expanding, resulting in continuous expansion of the tumor. Thus, the identification of mechanisms in which cancer cells regenerate themselves through a process called self-renewal is critical to our understanding of these diseases. It is becoming clear that self-renewal pathways in at least some of the cancer cells are disrupted, resulting in an expansion of tumorigenic cancer cells. To develop assays to identify these pathways, one must first identify which cancer cells in the tumor are capable of self-renewal.

Several genes that regulate normal stem cell self-renewal play a role in cancer. Recently, it has been shown that Wnt/ β -catenin signaling plays a pivotal role in both self-renewal of normal stem cells and malignant transformation [7, 17, 27]. The Wnt pathway (Fig. 2) was first discovered in mouse mammary tumor virus-induced breast cancer where deregulated expression of Wnt-1 due to proviral insertion resulted in mammary tumors [23, 32]. Subsequently, it has been shown that Wnt proteins play a central role in pattern formation. Wnt-1 belongs to a large family of highly hydrophobic secreted proteins that function by binding to their cognate receptor, a dimer of one of the members of the Frizzled family with low-density lipoprotein receptor-related protein 5 or 6, resulting in the activation of β -catenin [7]. In the absence of receptor activation, β -catenin is marked for degradation by a complex consisting of the adenomatous polyposis coli (APC), Axin and glycogen synthase kinase-3 β proteins [14, 30].

Wnt proteins are expressed in the bone marrow, and activation of Wnt/ β -catenin signaling by Wnt in vitro or by expression of a constitutively active β -catenin in vivo expands the pool of early progenitor cells and enriched normal transplantable hematopoietic stem cells in tissue culture and in vivo [27]. The inhibition of Wnt/ β -catenin by ectopic expression of Axin, which targets β -catenin for ubiquitination and degradation, leads to the inhibition of stem cell proliferation both in vitro and in vivo. Other studies suggest that the Wnt/ β -catenin pathway mediates stem or progenitor cell self-renewal in other tissues [32]. Higher levels of β -catenin are seen in keratinocytes with higher proliferative potential than those seen in keratinocytes with lower proliferative capacity [11]. Like their normal hematopoietic stem cell




Fig. 1 Flow cytometry of a breast cancer tumor. Pleural effusion cells were dissociated and stained with antibodies against CD44, CD24, and a panel of antibodies that recognize normal cells [2]. The $CD44^{+} CD24^{-/low}$ cancer cells, but not the majority of cancer cells with other phenotypes, had the exclusive ability to form tumors




Fig. 2 Wnt signaling cascades. In normal cells, E-cadherin and α -catenin link β -catenin to the cell membrane and the cytoskeleton. In the canonical Wnt pathway, Wnt ligands bind to Frizzled receptors utilizing the LRP 5/6 co-receptors. DKK1 is an extracellular inhibitor of Wnt signaling that blocks the LRP 5/6 co-receptors. Ligand-bound Frizzled receptors activate Dishevelled (*DSH*) by an unknown mechanism, inhibiting the glycogen synthase kinase-3 (*GSK-3*)/Axin/adenomatous polyposis coli (*APC*) complex that targets β -catenin for degradation via the ubiquitination pathway in the absence of Wnt signaling. β -Catenin translocates to the nucleus, where it interacts with TCF/LEF

transcription factors to initiate transcription of target genes. In the alternative pathway, the binding of Wnt ligands to Frizzled and LRP 5/6 co-receptors activates heterotrimeric G proteins (G). The ? indicates that disruption of the E-cadherin/ β -catenin complex may release β -catenin from the adhesion complex and allow it to activate the canonical Wnt-signaling pathway. Activation of G proteins results in the formation of inositol-1,4,5-triphosphate (IP_3), which triggers the release of intracellular Ca^{2+} . This triggers the activation of the Ca^{2+} -sensitive enzymes such as calmodulin-dependent protein kinase II (*CamKII*) and protein kinase C (*PKC*), which leads to cellular responses (*DAG* diacylglycerol)


counterparts, enforced expression of an activated β -catenin increased the ability of epidermal stem cells to self-renew and decreased their ability to differentiate. Mice that fail to express TCF-4, one of the transcription factors that is activated when bound to β -catenin, soon exhaust their undifferentiated crypt epithelial progenitor cells, further suggesting that Wnt signaling is involved in the self-renewal of epithelial stem cells. Normally, β -catenin is anchored to the cell surface and the cytoskeleton by E-cadherin and α -catenin (Fig. 2). In the fruit fly, disruption of this interaction leads to increased stem cell proliferation and expansion of the stem cell pool, probably secondary to loss of polar cell divisions [36].

The catenins have been implicated in cancer. Activation of β -catenin in colon cancer by inactivation of the protein degradation pathway, most frequently by mutation of APC, is common [31]. Expression of certain Wnt genes is elevated in some other epithelial cancers, suggesting that activation of β -catenin might be secondary to ligand activation in such cancers [3]. There is evidence that constitutive activation of the

Wnt/ β -catenin pathway may confer a stem/progenitor cell phenotype to colon cancer cells. Inhibition of β -catenin/TCF-4 in a colon cancer cell line induced the expression of the cell-cycle inhibitor p21^{cip-1}, and also induced the cells to stop proliferating and to acquire a more differentiated phenotype [33]. Enforced expression of the proto-oncogene *c-myc*, which is transcriptionally activated by β -catenin/TCF-4, inhibited the expression of p21^{cip-1} and allowed the colon cancer cells to proliferate when β -catenin/TCF-4 signaling was blocked, linking Wnt signaling to *c-myc* in the regulation of cell proliferation and differentiation.

The role of the mutations in the components of the canonical β -catenin pathway in breast cancer is less clear [9]. Unlike colon cancer, in breast cancer, mutations in APC or β -catenin resulting in stabilization of the protein are rare. In colon cancer, these mutations result in the accumulation of the β -catenin in the nucleus. Of the 24 breast cancer cell lines studied, only the DU 4475 cells contained a mutant APC gene [28]. Despite the rarity of mutations in APC or β -catenin, in some cases of breast cancer, the β -catenin is located in the nucleus rather than

Fig. 3 Effect of cancer treatments. When patients with solid tumors are treated, their tumors often shrink. Since the bulk of the tumor is not composed of the cancer stem cells, the therapies must eliminate the nontumorigenic cells. In the case of testicular cancer, even though platinum-based therapies do not always eliminate all of the cancer cells, the cancer stem cells are usually eliminated and most patients are cured. However, in most solid tumors, the therapies likely spare a significant number of the cancer stem cells and the residual cancer stem cells regenerate the tumor



the outer membrane. In these cancers, *c-myc* and cyclin D1, downstream targets of β -catenin, are overexpressed and this is associated with poor prognosis. Studies using cell lines suggest that in some cancer cell lines the β -catenin pathway is activated by autocrine secretion of Wnts. In these cells, growth is inhibited using the soluble Wnt inhibitor DKK1 and the growth of some breast cancer cell lines is inhibited.

division and proliferation would help them to avoid chemotherapy and radiation. Certain antigens currently targeted by biologic therapies may not be expressed on cell surfaces until the cells are more mature. Current therapy may be good at producing initial tumor burden reduction, but if the cancer stem cells are spared, relapse is inevitable (Fig. 3).

Implications of cancer stem cells

An axiom in the treatment of tumors is that remission is, in general, more difficult to achieve with each relapse. Obtaining initial remission as dictated by current surveillance methods only results in the patient succumbing to disease in relapse. Metastasis is also a difficult hurdle to cross in many clinical settings. In most cancers, the presence of metastasis at diagnosis dictates more aggressive therapies, and lower disease-free survival rates, as seen in sarcomas and neuroblastomas. In the cancer stem cell model, it is the tumorigenic cancer stem cell that escapes chemotherapy and metastasizes to a new location to cause distant tumor recurrence. Our current approaches to the cure are dependent on a few qualities that tumors exhibit. Surgical approaches are successful in tumors in which metastasis is not an issue and where the tumor can be removed en bloc. Many chemotherapeutic drugs and radiation treatments depend on cells that divide and proliferate at high rates. Antibody therapy is dependent on the presence of an effective antigen. Under the cancer stem cell model, it becomes clear that present therapy would not be effective at targeting cancer stem cells. If cancer stem cells exhibit qualities of other stem cells, their low rates of

Conclusions

Cancers arise in tissues that contain a stem cell population. Since stem cells are often the longest-lived cells in an organ, mutations leading to cancer often accumulate in the stem cell pool. Tumors of many, if not all, organs contain a cancer stem cell pool that has the exclusive ability to drive tumor growth. These cells may or may not be derived from normal stem cells. Regardless of the cell of origin, the cancer stem cells rely on self-renewal pathways present in normal stem cells to maintain themselves and expand. Clearly, there are mutations in the self-renewal pathways in cancer stem cells. Targeting these aberrant self-renewal pathways in cancer cells may result in more effective cancer therapies.

References

1. Akashi K, Weissman IL (2001) In: Zon LI (ed) Developmental biology of hematopoiesis. Oxford University Press, New York
2. Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF (2003) Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci USA* 100:3983–3988
3. Bafico A, Liu G, Goldin L, Harris V, Aaronson SA (2004) An autocrine mechanism for constitutive Wnt pathway activation in human cancer cells. *Cancer Cell* 6:497–506

4. Bergsagel DE, Valeriote FA (1968) Growth characteristics of a mouse plasma cell tumor. *Cancer Res* 28:2187–2196
5. Brown JM (1997) NCI's anticancer drug screening program may not be selecting for clinically active compounds. *Oncol Res* 9:213–215
6. Bruce WR, Gaag H (1963) A quantitative assay for the number of murine lymphoma cells capable of proliferation in vivo. *Nature* 199:79–80
7. Cadigan KM, Nusse R (1997) Wnt signaling: a common theme in animal development. *Genes Dev* 11:3286–3305
8. Dorrell C, Gan OI, Pereira DS, Hawley RG, Dick JE (2000) Expansion of human cord blood CD34(+)CD38(–) cells in ex vivo culture during retroviral transduction without a corresponding increase in SCID repopulating cell (SRC) frequency: dissociation of SRC phenotype and function. *Blood* 95:102–110
9. Ethier SP, Mahacek ML, Gullick WJ, Frank TS, Weber BL (1993) Differential isolation of normal luminal mammary epithelial cells and breast cancer cells from primary and metastatic sites using selective media. *Cancer Res* 53:627–635
10. Furley AJ, Reeves BR, Mizutani S, Altass LJ, Watt SM, Jacob MC, van den Elsen P, Terhorst C, Greaves MF (1986) Divergent molecular phenotypes of KG1 and KG1a myeloid cell lines. *Blood* 68:1101–1107
11. Gat U, DasGupta R, Degenstein L, Fuchs E (1998) De novo hair follicle morphogenesis and hair tumors in mice expressing a truncated beta-catenin in skin. *Cell* 95:605–614
12. Hamburger AW, Salmon SE (1977) Primary bioassay of human tumor stem cells. *Science* 197:461–463
13. Hanahan D, Weinberg RA (2000) The hallmarks of cancer. *Cell* 100:57–70
14. Hedgepeth CM, Deardorff MA, Rankin K, Klein PS (1999) Regulation of glycogen synthase kinase 3beta and downstream Wnt signaling by Axin. *Mol Cell Biol* 19:7147–7157
15. Hoffman RM (1999) Orthotopic metastatic mouse models for anticancer drug discovery and evaluation: a bridge to the clinic. *Invest New Drugs* 17:343–359
16. Ikeda H, Kanakura Y, Furitsu T, Kitayama H, Sugahara H, Nishiura T, Karasuno T, Tomiyama Y, Yamatodani A, Kanayama Y, Matsuzawa Y (1993) Changes in phenotype and proliferative potential of human acute myeloblastic leukemia cells in culture with stem cell factor. *Exp Hematol* 21:1686–1694
17. Korinek V, Barker N, Moerer P, van Donselaar E, Huls G, Peters PJ, Clevers H (1998) Depletion of epithelial stem-cell compartments in the small intestine of mice lacking Tcf-4. *Nat Genet* 19:379–383
18. Krasna L, Dudorkinova D, Vedralova J, Vesely P, Pokorna E, Kudlackova I, Chaloupkova A, Petruzalka L, Danes J, Matouskova E (2002) Large expansion of morphologically heterogeneous mammary epithelial cells, including the luminal phenotype, from human breast tumours. *Breast Cancer Res Treat* 71:219–235
19. Lagasse E, Weissman IL (1994) bcl-2 inhibits apoptosis of neutrophils but not their engulfment by macrophages. *J Exp Med* 179:1047–1052
20. Leglise MC, Dent GA, Ayscue LH, Ross DW (1988) Leukemic cell maturation: phenotypic variability and oncogene expression in HL60 cells: a review. *Blood Cells* 13:319–337
21. Morrison SJ, Qian D, Jerabek L, Thiel BA, Park IK, Ford PS, Kiel MJ, Schork NJ, Weissman IL, Clarke MF (2002) A genetic determinant that specifically regulates the frequency of hematopoietic stem cells. *J Immunol* 168:635–642
22. Muller-Sieburg CE, Cho RH, Sieburg HB, Kupriyanov S, Riblet R (2000) Genetic control of hematopoietic stem cell frequency in mice is mostly cell autonomous. *Blood* 95:2446–2448
23. Nusse R, Brown A, Papkoff J, Scambler P, Shackleford G, McMahon A, Moon R, Varmus H (1991) A new nomenclature for int-1 and related genes: the Wnt gene family. *Cell* 64:231
24. Park CH, Bergsagel DE, McCulloch EA (1971) Mouse myeloma tumor stem cells: a primary cell culture assay. *J Natl Cancer Inst* 46:411–422
25. Phillips RL, Reinhart AJ, Van Zant G (1992) Genetic control of murine hematopoietic stem cell pool sizes and cycling kinetics. *Proc Natl Acad Sci USA* 89:11607–11611
26. Reya T, Morrison SJ, Clarke MF, Weissman IL (2001) Stem cells, cancer, and cancer stem cells. *Nature* 414:105–111
27. Reya T, Duncan AW, Ailles L, Domen J, Scherer DC, Willert K, Hintz L, Nusse R, Weissman IL (2003) A role for Wnt signalling in self-renewal of haematopoietic stem cells. *Nature* 423:409–414
28. Schloschauer PW, Brown SA, Eisinger K, Yan Q, Guglielminetti ER, Parsons R, Ellenson LH, Kitajewski J (2000) APC truncation and increased beta-catenin levels in a human breast cancer cell line. *Carcinogenesis* 21:1453–1456
29. Southam C, Brunschwig A (1961) Quantitative studies of autotransplantation of human cancer. *Cancer* 14:971–978
30. Spink KE, Polakis P, Weis WI (2000) Structural basis of the Axin-adenomatous polyposis coli interaction. *Embo J* 19:2270–2279
31. Taipale J, Beachy PA (2001) The hedgehog and Wnt signalling pathways in cancer. *Nature* 411:349–354
32. Tsukamoto AS, Grosschedl R, Guzman RC, Parslow T, Varmus HE (1988) Expression of the int-1 gene in transgenic mice is associated with mammary gland hyperplasia and adenocarcinomas in male and female mice. *Cell* 55:619–625
33. van de Wetering M, Sancho E, Verweij C, de Lau W, Oving I, Hurlstone A, van der Horn K, Battle E, Coudreuse D, Haramis AP, Tjon-Pon-Fong M, Moerer P, van den Born M, Soete G, Pals S, Eilers M, Medema R, Clevers H (2002) The beta-catenin/TCF-4 complex imposes a crypt progenitor phenotype on colorectal cancer cells. *Cell* 111:241–250
34. Weidmann E, Brieger J, Karakas T, Maurer U, Pascheberg U, Hoelzer D, Mitrou PS, Bergmann L (1997) Establishment and characterization of a new, factor-independent acute myeloid leukemia line designated Ei501. *Leukemia* 11:709–713
35. Wodinsky I, Swiniarski J, Kensler CJ (1967) Spleen colony studies of leukemia L1210.I. Growth kinetics of lymphocytic L1210 cells in vivo as determined by spleen colony assay. *Cancer Chemother Rep* 51:415–421
36. Yamashita YM, Jones DL, Fuller MT (2003) Orientation of asymmetric stem cell division by the APC tumor suppressor and centrosome. *Science* 301:1547–1550