

## REVIEW

# Human haematopoietic stem cell development: from the embryo to the dish

Andrejs Ivanovs<sup>1,2</sup>, Stanislav Rybtsov<sup>1</sup>, Elizabeth S. Ng<sup>3,4</sup>, Edouard G. Stanley<sup>3,4,5</sup>, Andrew G. Elefanty<sup>3,4,5,\*</sup> and Alexander Medvinsky<sup>1,\*</sup>

## ABSTRACT

Haematopoietic stem cells (HSCs) emerge during embryogenesis and give rise to the adult haematopoietic system. Understanding how early haematopoietic development occurs is of fundamental importance for basic biology and medical sciences, but our knowledge is still limited compared with what we know of adult HSCs and their microenvironment. This is particularly true for human haematopoiesis, and is reflected in our current inability to recapitulate the development of HSCs from pluripotent stem cells *in vitro*. In this Review, we discuss what is known of human haematopoietic development: the anatomical sites at which it occurs, the different temporal waves of haematopoiesis, the emergence of the first HSCs and the signalling landscape of the haematopoietic niche. We also discuss the extent to which *in vitro* differentiation of human pluripotent stem cells recapitulates bona fide human developmental haematopoiesis, and outline some future directions in the field.

**KEY WORDS:** HSC, Human, Embryo, hPSC, Reprogramming

## Introduction

In mammals, the first transient waves of blood cells arise in the yolk sac (extra-embryonic haematopoiesis) and serve the immediate needs of the growing embryo (Silver and Palis, 1997). They include the most primitive wave, which consists mainly of large nucleated erythrocytes that is rapidly followed by a second wave of erythromyeloid progenitors (EMPs) and lymphoid progenitors that transiently seed the foetal liver (Böiers et al., 2013; McGrath et al., 2015). The third wave, which includes self-renewing haematopoietic stem cells (HSCs) that give rise to the permanent adult haematopoietic system, emerges later inside the body of the embryo (intra-embryonic haematopoiesis) in the aorta-gonad-mesonephros (AGM) region, which is evolutionarily conserved in many vertebrates (Medvinsky et al., 1993; Medvinsky and Dzierzak, 1996). The process of HSC development in vertebrates is driven by largely conserved, although not entirely identical, molecular mechanisms (Medvinsky et al., 2011; Ciau-Uitz et al., 2016). The endothelial origin of haematopoietic progenitors and HSCs has been established by their emergence from the aortic endothelial layer (Garcia-Porrero et al., 1995; Tavian et al., 1996;

Jaffredo et al., 1998), lineage tracing *in vivo* using genetic labelling (Zovein et al., 2008; Chen et al., 2009; Bertrand et al., 2010; Kissa and Herbomel, 2010) and observations of embryonic stem cells (ESCs) differentiating *in vitro* (Eilken et al., 2009; Lancrin et al., 2009). HSCs and non-self-renewing haematopoietic progenitor cells that emerge in the AGM region are organised within intra-aortic haematopoietic clusters (IAHCs). These bud predominantly from the endothelial floor of the dorsal aorta, which accordingly co-express endothelial and haematopoietic markers. The search for markers that would accurately identify developing HSCs and separate them from non-self-renewing progenitors is an important goal towards understanding the mechanisms that underlie HSC development (Rybtsov et al., 2014; Zhou et al., 2016). Important events that occur downstream of HSC emergence include colonisation and expansion of HSCs in the foetal liver and subsequent lodging in the adult bone marrow.

Our current understanding of the mechanisms that drive HSC development has come from analysis of various model organisms. We do not intend here to provide detailed overview of non-human models, as this extensive topic has been reviewed recently by us and by others (Medvinsky et al., 2011; Kim et al., 2014; Ciau-Uitz et al., 2016; Kauts et al., 2016; Robertson et al., 2016; Crisan and Dzierzak, 2016). Owing to the high accessibility of model organisms for experimentation and the genetic variability and rarity of human embryonic material, the analysis of human haematopoietic development has lagged behind these well-established experimental models and for a long time was limited to immunohistological and *in vitro* studies. More recently, the development of highly efficient xenograft mouse models has allowed human HSCs to be tested functionally (Shultz et al., 2012). Despite the discovery of major commonalities between mouse and human HSC development, significant differences remain that need to be characterised in order to understand human HSC development.

Attempts to recapitulate haematopoiesis development via directed differentiation of pluripotent stem cells (PSCs) – both ESCs and induced PSCs (iPSCs) – have provided a valuable source of information, although current protocols appear to replicate extra-embryonic, yolk sac-like, rather than intra-embryonic, adult-like haematopoiesis. Indeed, it may be for this reason that efforts to generate bona fide HSCs from PSCs *in vitro* have not been successful, undermining expectations for the generation of HSCs for clinical needs. This, in turn, has stimulated the quest for alternative methodologies, such as direct cell reprogramming (Riddell et al., 2014; Sandler et al., 2014). In this Review, we first discuss *in vivo* human haematopoietic development, focussing on the different anatomical sites where haematopoiesis takes place, as well as the molecular and functional characterisation of human haematopoietic stem and progenitor cells as they emerge in a spatiotemporal manner. Based on what is known regarding the development of the human haematopoietic system, we consider the

<sup>1</sup>Institute for Stem Cell Research, MRC Centre for Regenerative Medicine, University of Edinburgh, Edinburgh EH16 4UU, UK. <sup>2</sup>Institute of Anatomy and Anthropology, Riga Stradiņš University, Riga LV-1007, Latvia. <sup>3</sup>Murdoch Childrens Research Institute, The Royal Children's Hospital, Parkville, Victoria 3052, Australia. <sup>4</sup>Department of Anatomy and Developmental Biology, Faculty of Medicine, Nursing and Health Sciences, Monash University, Clayton, Victoria 3800, Australia. <sup>5</sup>Department of Paediatrics, Faculty of Medicine, Dentistry and Health Sciences, University of Melbourne, Parkville, Victoria 3052, Australia.

\*Authors for correspondence (a.medvinsky@ed.ac.uk; andrew.elefanty@mcri.edu.au)

extent to which attempts to generate HSCs *in vitro* recapitulate *in vivo* development, and discuss the prospect of generating true fully functional HSCs in the dish – cells that would be suitable for transplantations in clinical settings. The generation of HSCs *in vitro* in the absence of genetic modifications would represent the strongest indication that the key basic mechanisms that underpin HSC development in the human embryo have become largely understood.

### The anatomy of haematopoietic development

In the mouse, developmental stages can be distinguished by morphological changes occurring over less than a day. Human embryo development, however, takes a significantly longer time. Stages of human development, known as Carnegie stages (CS), are defined by external morphological characteristics and normally cover several days each (O’Rahilly and Muller, 1987). Staging according to the Carnegie classification is more accurate than using gestational age or crown-rump length, which is important when analysing human HSC development (Ivanovs et al., 2013). For consistency, where days post-conception (dpc) were used in original publications, corresponding CS are presented in parallel in this Review. Despite differences in developmental timescales, substantial parallels between mouse and human haematopoietic development can readily be observed (Fig. 1).

#### The yolk sac

The first wave of human haematopoiesis originates in the yolk sac, a structure that develops differently in human and mouse. Whereas in the mouse conceptus the yolk sac surrounds the body of the embryo, in the human it develops in front of the embryo body in a balloon-like form (Fig. 2A). Tight groups of ‘mesenchymal’ cells, of mesoderm origin, that are adjacent to the endoderm, differentiate into haematopoietic cells surrounded by endothelial cells, which later remodel to form the yolk sac vascular plexus. Large primitive nucleated erythrocytes represent the major haematopoietic output from the yolk sac at CS 7–8 (16–18.5 dpc), with the occasional presence of primitive macrophages and megakaryocytes (Bloom and Bartelmez, 1940; Fukuda, 1973; Luckett, 1978). By CS 10 (21–22 dpc), the first primitive erythroblasts can be observed inside the cardiac cavity, marking the onset of blood circulation, followed by appearance of the first CD45<sup>+</sup> (PTPRC<sup>+</sup>) cells (Tavian et al., 1999), similar to analyses in the mouse (Ghiaur et al., 2008). Although a yolk sac EMP wave similar to the mouse has not been formally characterised during human development, early studies indicate that this wave may emerge in the human yolk sac during CS 13–15 (28–35 dpc) (Migliaccio et al., 1986).

#### The AGM region

The appearance of IAHCs on the ventral wall of the human dorsal aorta in the AGM region, reported early last century (Minot, 1912), marks the onset of the intra-embryonic, permanent adult haematopoietic wave in the vertebrate embryo. Spatiotemporal analysis revealed that IAHCs emerge at CS 13 (27 dpc), exclusively at the floor of the embryonic dorsal aorta and disappear by CS 17 (39–42 dpc) (Tavian et al., 1996, 1999). By contrast, in the mouse some IAHCs are also observed in the aortic roof (Taoudi and Medvinsky, 2007; Taylor et al., 2010; Yokomizo and Dzierzak, 2010). In the human embryo, IAHC formation covers the pre-umbilical area of the floor of the dorsal aorta and penetrates the entry to the vitelline artery (Fig. 2B,C). Early mesoderm and the endothelial compartment in early post-gastrulation embryos is marked by KDR (also known as FLK1) expression (Cortés et al.,

1999). The endothelial lining of the human dorsal aorta upregulates CD34, which later marks adult HSCs, from CS 9 (19 dpc) (Tavian et al., 1996, 1999, 2001; Oberlin et al., 2002). By CS 13 (27 dpc), the first CD34<sup>+</sup>CD45<sup>+</sup> cells emerge in the pre-umbilical region of the dorsal aorta, and their number reaches several hundred by CS 15 (33 dpc).


Although it is broadly accepted that IAHCs are formed from mesodermal precursors through endothelial intermediates, in the absence of direct tracking, various scenarios of IAHC formation can be considered (Medvinsky et al., 2011) (see also ‘The endothelial origin of human haematopoiesis’, below). Developing human IAHCs sometimes appear to penetrate through the aortic endothelial lining (Tavian et al., 1999). This could reflect a number of scenarios, including ‘excessive’ endothelial-to-haematopoietic transition (EHT) activity *in situ*, transendothelial cell migration towards the aortic lumen into the bloodstream (Bertrand et al., 2005; Rybtsov et al., 2011), or migration of EHT-undergoing cells in the opposite direction, towards the venous system, as described in zebrafish (Bertrand et al., 2010; Kissa and Herbolme, 2010). Expression of angiotensin converting enzyme (ACE), which labels human foetal liver HSCs (Jokubaitis et al., 2008), identifies scattered ACE<sup>+</sup>CD34<sup>−</sup> cells beneath the dorsal aorta (Sinka et al., 2012). These have been postulated to represent IAHC precursors, which upregulate CD34 and after integration into the CD34<sup>+</sup> endothelial lining form ACE<sup>+</sup>CD34<sup>+</sup> IAHCs (Sinka et al., 2012). Identification of the exact migration pathways during HSC specification in the mammalian AGM region remains one of the least explored issues in the field.

#### The liver

The liver rudiment emerges as a diverticulum from the floor of the embryonic gut at early CS 10 (21 dpc). From late CS 10 (22 dpc), the liver rudiment contains primitive yolk sac-derived erythrocytes and CD45<sup>+</sup> cells, likely of monocytic/macrophage lineage. From CS 13 (27–29 dpc), the liver is seeded by growing numbers of CD34<sup>+</sup>CD45<sup>+</sup> cells (Tavian et al., 1999), which likely represent yolk sac-derived cells similar to mouse EMPs (McGrath et al., 2015). Although direct experimental evidence is lacking, these might be the cells that progressively replace primitive erythroblasts in the human embryo bloodstream during liver colonisation from CS 13–14 (30–33 dpc) onwards (Migliaccio et al., 1986). Concurrent emergence of IAHCs and CD34<sup>+</sup>CD45<sup>+</sup> cells in the liver at CS 13 suggests that some of these cells also represent colonisation of the foetal liver by AGM-derived cells, despite a delay in the appearance of HSCs in the liver (see ‘Spatiotemporal HSC development in the early human embryo’, below). The liver remains an important niche for haematopoietic differentiation and HSC expansion until birth. Although some analysis of the developing liver as the haematopoietic niche was conducted on the mouse, this issue in human remains a largely unexplored territory.

#### The placenta

In the mouse placenta, definitive HSCs are detected at the same developmental age as in the AGM region. For this reason, the placenta is considered to be one of the HSC sources in the mouse embryo (Gekas et al., 2005; Ottersbach and Dzierzak, 2005). At least from week 5–6 of development, the human placenta contains high numbers of CD34<sup>+</sup> cells, represented by immature CD34<sup>+</sup>CD45<sup>lo</sup> cells that lack CD38 and contain high colony-forming units-culture (CFU-C) numbers, and CD34<sup>+</sup>CD45<sup>lo</sup> cells committed to erythroid and myeloid differentiation containing fewer CFU-Cs (Barcena et al., 2009). At this stage, the placenta is also a site of extensive erythroid maturation: placental villi are



**Fig. 1. Chronology of human haematopoietic development.** The first human haematopoietic cells, primitive erythroid cells and monocytes/macrophages are produced in the yolk sac during CS 7 and 8 (16–18.5 days). With the onset of cardiac contractions at CS 10 (day 21) and blood circulation, yolk sac-derived haematopoietic cells are disseminated throughout the developing embryo. CD34<sup>+</sup>CD45<sup>+</sup> intra-aortic haematopoietic clusters appear in the vitelline artery and on the ventral wall of the dorsal aorta at CS 13 (day 27). The clusters disappear by CS 16 (35–38 days). HSCs in the AGM region persist from CS 14 (day 30) until at least CS 17 (day 42; latest stage tested). HSC activity in the AGM region precedes that in the yolk sac (CS 16; 35–38 days), liver (CS 17; 39–42 days) and placenta (ca. 63 days). The haematopoietic lineage derives from a mesodermal precursor (grey) through the intermediate of the haematogenic endothelium before expanding. For the sake of comparison, the step-wise emergence of mouse HSCs through immature haematopoietic precursors (pro-HSC, pre-HSC I, pre-HSC II) is shown at the bottom of the figure. The existence of similar precursors in human haematopoietic development has not been functionally shown, which is indicated by a question mark for human pro/pre-HSC. Red, bona fide HSCs; light red, haematopoietic lineages which may or may not be related to HSC development; yellow, yolk sac haematopoietic differentiation. Fading of coloured bubbles to blue represents extinction of the process. White striped lines represent a change in time scale (omission of several days for the mouse or weeks for the human).


enriched for primitive erythrocytes that express embryonic  $\zeta$ -globin in association with macrophages, which may facilitate their enucleation (Van Handel et al., 2010). Although CD34<sup>+</sup> cells appear in the human placenta as early as at week 5 of gestation, true HSCs can be detected there only after week 9 of gestation or even later, as determined by xenotransplantation into immunocompromised mice (Robin et al., 2009; Muench et al., 2017). Although the CD34<sup>+</sup>CD45<sup>lo</sup> population that contains HSCs

is embedded within a vimentin-positive stromal environment or rarely in close association with blood vessels (Muench et al., 2017), localisation of the exact HSC fraction and their regulatory microenvironment in the human placenta is currently unavailable.

#### The bone marrow

The development of the definitive bone marrow niche is closely linked to the invasion of cartilaginous bone by blood vessels and





**Fig. 2. Human embryonic haematopoietic tissues.** (A) The location of the AGM region, yolk sac, liver, vitelline and umbilical arteries and placenta in a CS 15 human embryo. The location of the amnion, eye, and upper and lower limbs are also indicated for reference. Note that HSCs at this stage are localised to the AGM region and will appear in the yolk sac and liver slightly later. (B) Human IAH development and the stage-specific spatial localisation of CD34<sup>+</sup>CD45<sup>+</sup> cell clusters on the ventral wall of the dorsal aorta [indicated in red; modified from Tavian and Peault (2005)]. (C) VE-cadherin<sup>+</sup>CD45<sup>+</sup>RUNX1<sup>+</sup> human IAHs on the ventral wall of the dorsal aorta of a CS 16 human embryo. Whole-mount antibody staining (left) and sagittal confocal section of the boxed area (right). The largest clusters are usually localised close to the entry of the vitelline artery. Scale bars: 1 mm in A; 0.05 mm in C.

bone ossification. Vascular invasion facilitates seeding of bone marrow with haematopoietic progenitors and HSCs. Bone marrow formation arbitrarily marks the end of the human embryonic period (CS 23; 56 dpc) (O’Rahilly and Muller, 1987). The initial CD34<sup>+</sup>CD45<sup>+</sup> haematopoietic cells that invade the cartilaginous bone include mostly CD68<sup>+</sup> monocytes/macrophages possibly participating in chondrolysis. This is followed by colonisation with CD34<sup>+</sup>CD45<sup>+</sup> progenitors and HSCs (Charbord et al., 1996), although the time of onset of HSC activity in the human bone marrow remains unknown.

#### Understanding haematopoietic development through *in vitro* approaches

A semi-solid *in vitro* (methylcellulose) assay allows an assessment of the presence of CD34<sup>+</sup> CFU-Cs in the early human embryo (Tavian et al., 1999). However, this assay does not reveal the potential of embryonic tissues to generate CFU-Cs or HSCs. In the mouse, this issue was addressed by pre-culturing embryonic tissues as explants to allow their further development *ex vivo* before assessing their haematopoietic potential (Cumano et al., 1996; Medvinsky and Dzierzak, 1996). This approach was adopted for human studies and revealed that the human yolk sac and para-aortic splanchnopleura (P-Sp; the precursor of the AGM region) isolated prior to the onset of circulation at CS 10 (21 dpc; to exclude cross-seeding) possess differing haematopoietic potentials (Tavian et al., 2001). The yolk sac could generate only myeloid and natural killer (NK) cells, whereas the P-Sp showed a broader spectrum of haematopoietic differentiation including lymphoid B- and T-cell

lineages starting from CS 11 (24 dpc), suggesting that it could give rise to multipotent haematopoietic progenitors *in vivo* (Tavian et al., 2001). A subsequent study revealed that only the dorsal aorta and not the yolk sac contained so-called long-term cobblestone area-forming cells, which in bone marrow correlate with the presence of HSCs (Oberlin et al., 2002). Although these results are consistent with analyses in the mouse embryo, the human lympho-myeloid precursors revealed in culture may not be equal to true HSCs. Identification of the very few HSCs in the early human embryo required development of a robust *in vivo* long-term repopulation assay, which became possible only with the later advent of highly receptive xenograft mouse models.

#### The endothelial origin of human haematopoiesis

Various non-human experimental models have shown that during development haematopoietic cells emerge from the embryonic endothelium (Jaffredo et al., 2000; de Bruijn et al., 2002; Bertrand et al., 2010; Kissa and Herbomel, 2010; Chen et al., 2011). KDR<sup>+</sup>CD34<sup>+</sup> cells observed in the early human embryo mark mesodermal precursors that are likely to develop into endothelial precursors (Cortés et al., 1999). CD34, which initially marks human aortic endothelium, at CS 13 (27 dpc) has also been shown to label IAHs (Tavian et al., 1999). In a more recent study, CD34<sup>+</sup>CD45<sup>+</sup> populations purified from CS 13–18 (28–44 dpc) dorsal aortas and vitelline arteries were shown to give rise to myeloid and lymphoid populations after co-culture with stromal cells (Oberlin et al., 2002). This activity of the CD34<sup>+</sup>CD45<sup>+</sup> population (defined as haematogenic endothelium) temporally correlated with IAH

formation, supporting the idea that clonogenic haematopoietic progenitors reside within these structures. The CD34<sup>+</sup>CD45<sup>−</sup> population, however, is heterogeneous and may contain not only endothelial cells. Indeed, the endothelial-to-haematopoietic transition (EHT) is a multistep maturation process, which includes cell intermediates fully committed to the haematopoietic fate (Taoudi et al., 2008; Medvinsky et al., 2011; Rybtsov et al., 2014). In the mouse embryo and differentiating ESCs, the earliest haematopoietic progenitors express the haematopoietic marker CD41 (ITGA2B), and only later become CD45<sup>+</sup> (Ferkowicz et al., 2003; Mikkola et al., 2003; Eilken et al., 2009; Lancrin et al., 2009). Similarly, the developing mouse HSC lineage sequentially upregulates CD41, CD43 and finally CD45, which together define the different stages of HSC maturation (Rybtsov et al., 2014). Importantly, although these cell intermediates express perhaps the most specific endothelial marker, VE-cadherin (cadherin 5), they are devoid of endothelial activity and are fully committed to the haematopoietic fate. During human ESC differentiation *in vitro*, CD43 is the first marker of haematopoietic specification (Vodyanik et al., 2006). Therefore, apart from endothelial cells, the human AGM-derived CD34<sup>+</sup>CD45<sup>−</sup> population might also include haematopoietic progenitors, which could be responsible for haematopoietic activity in culture (Oberlin et al., 2002). Although some molecular mechanisms underlying EHT transition have been described (Ciau-Uitz et al., 2016), exactly how haematopoietic progenitors and HSCs emerge from the embryonic endothelium, especially in the human, remains an issue for further investigation.

### Spatiotemporal HSC development in the early human embryo

Human HSC development has been functionally evaluated through transplantation into immunodeficient mice (see Box 1), which represents a more stringent, long-term and robust assessment than *in vitro* assays. All haematopoietic tissues known from mouse studies, such as the AGM region, yolk sac, liver, umbilical cord and placenta (Fig. 2A) have been assayed (Ivanovs et al., 2011). Short tandem repeat analysis allowed for the discrimination between embryonic and maternal cells, which might contaminate samples, especially in

placenta transplants. Examination of CS 12–17 (25–42 dpc) embryos revealed the presence of HSCs predominantly in the AGM region. Out of 11 experiments in which multipotent HSCs were detected in the AGM region, nine showed the presence of HSCs solely in that location. The first HSCs in the AGM region were detected as early as at CS 14 (30–32 dpc) and persisted until at least CS 17 (39–42 dpc). In the yolk sac, HSCs were detected only after CS 16 (35–38 dpc), approximately 5 days later than in the AGM region. In general, repopulation with yolk sac cells was significantly rarer than with AGM region cells. A robust presence of HSCs in the liver is generally detected later, from week 7–8 and the most potent subset of liver HSCs still express VE-cadherin, reflecting their endothelial origin (Oberlin et al., 2010; Ivanovs et al., 2011).

Although the appearance of HSCs in the mouse AGM region, yolk sac, umbilical cord, liver and placenta occur almost concurrently (Müller et al., 1994; de Bruijn et al., 2000; Kumaravelu et al., 2002; Gekas et al., 2005; Ottersbach and Dzierzak, 2005), their appearance in different locations in human embryo is temporally resolved, owing to a more protracted developmental period. Whereas in the mouse HSCs develop in umbilical cord and placenta prior to liver colonisation (de Bruijn et al., 2000; Gekas et al., 2005; Ottersbach and Dzierzak, 2005), human umbilical cord and placenta at analogous stages (CS 14–17; 30–42 dpc) lack HSCs (Ivanovs et al., 2011). Only from week 9 of development are HSCs reliably detected in the human placenta (Robin et al., 2009), suggesting that the human placenta is a secondary site for HSC development.


The aforementioned studies emphasise the importance of functional *in vivo* assays for the analysis of HSCs. Although immunohistological and *in vitro* studies suggested the appearance of HSCs in the CS 13 (27–29 dpc) liver (Tavian et al., 1999), transplantable HSCs appear there only after CS 17 (39–42 dpc) (Ivanovs et al., 2011). Similarly, despite the presence of cells with an HSC phenotype and CFU-Cs, week 5–6 human placentas lack definitive HSCs (Barcena et al., 2009; Robin et al., 2009; Ivanovs et al., 2011).

### The exceptional regenerative potential of the first human HSCs

Transplantation of AGM cells after CS 14 (30–32 dpc) into immunodeficient mice results in long-term multilineage haematopoietic engraftment, confirming the presence of definitive HSCs (Ivanovs et al., 2011). As in the mouse, very few – normally only one – HSCs emerge in the human AGM region, as transplantation of cells from one human AGM into several recipients usually results in one haematopoietically reconstituted mouse (Fig. 3). Repopulation kinetics of human AGM region-derived HSCs are significantly slower than their mouse counterparts. By month 3 post-transplantation, haematopoietic chimerism reaches approximately 1% of total leukocyte count in the recipient blood. However, subsequent growth of human haematopoietic cells results in up to 90% chimerism by month 8 in recipient peripheral blood and bone marrow, which is accompanied by extensive generation of large numbers (more than 300) of fully functional daughter HSCs, as determined by re-transplantation into secondary recipients (Fig. 3) (Ivanovs et al., 2011). For comparison, this significant production of functional daughter HSCs exceeds the number required to prevent bone marrow failure in patients after myeloablation (Catlin et al., 2011). Computational analysis of the repopulation dynamics suggested that adult human HSCs replicate every 40 weeks (Catlin et al., 2011). However, amplification of human AGM-derived HSCs from 1 to 300 in the recipient mouse over 6 months requires

#### Box 1. The history of the (NOD/SCID/Gamma) NSG mouse line for haematopoietic transplantation assays

In the late 1980s, introduction of the antiprogesterone compound mifepristone for elective termination of early pregnancies (Vervest and Haspels, 1985) made early intact human embryos available for research purposes. At about the same time, severe combined immunodeficiency (SCID) mice, which could accept human haematopoietic grafts, were generated (McCune et al., 1988). These two factors largely defined progress in functional analysis of human HSCs. As SCID mice often immunoreject xenografts, they were backcrossed to non-obese diabetic (NOD) mice, which are deficient in functional T, B and antigen-presenting cells. For a long time, NOD/SCID recipient mice (Shultz et al., 1995) were widely used to study human haematopoiesis *in vivo*. However, owing to residual NK cell activity, haematopoietic engraftment of human HSCs is attenuated, which especially affected experiments with low HSC numbers. This, together with the short lifespan of these mice, diminishes the utility of the model. To address this limitation, NOD/SCID/Gamma (NSG) mice were engineered by introducing an additional mutation in the IL2 receptor  $\gamma$  chain. These mice, which lack NK cells, live for longer and support high-level human HSC engraftment (Shultz et al., 2005). The NSG mice later permitted transplantation analysis of the first human HSCs emerging in low numbers in the AGM region (Ivanovs et al., 2014, 2011).



**Fig. 3. Quantitative analysis of the regenerative potential of human AGM region-derived HSCs.** A cell suspension of dissociated AGM region tissue repopulated the haematopoietic system of just one primary recipient NSG mouse, indicating the presence of only one HSC in this tissue. Successful engraftment and thus the presence of a bona fide HSC is indicated by a red oval; non-HSC cells are indicated by black ovals. Six months after transplantation, the bone marrow (BM) of the primary recipient was transplanted in serial dilutions into secondary recipients. Repopulation of all recipients indicated that the original AGM-derived HSC generated no fewer than 300 daughter HSCs (Ivanovs et al., 2011).

replication at least every 2–4 weeks, because transplanted HSCs also generate differentiated blood cells. The proliferation of the first HSCs *in situ* (in the embryo) may be even higher, given massive expansion of HSCs in the foetal liver.

Achieving similar levels of immunodeficient mouse engraftment with human umbilical cord blood (UCB) HSCs requires considerably higher HSC doses, with 10–20 human UCB HSCs giving approximately 40% repopulation by 8 months post-transplantation (Notta et al., 2011). Re-transplantation of bone marrow from recipients reconstituted with low numbers of UCB-derived HSCs is inefficient and gives either low or no repopulation. The unprecedented superior self-renewal capacity of AGM region-derived HSCs is in line with observations that developmentally younger foetal liver HSCs are more potent than HSCs from adult bone marrow (Rebel et al., 1996; Holyoake et al., 1999).

### Localisation and phenotype of the first human HSCs

In mice, IAHCs are localised predominantly to the ventral domain of the dorsal aorta, correlating with the presence of functional HSCs (Taoudi and Medvinsky, 2007). Similarly, the ventral domain of the human AGM region is enriched for cells with an HSC immunophenotype and function in transplantation studies, suggesting that human HSCs also localised to IAHCs (Ivanovs

et al., 2014). However, although IAHCs disappear by CS 17 (35–38 dpc), HSCs are still detectable at this stage indicating that functional HSCs in the human AGM region do not entirely correlate with the presence of IAHCs (Tavian et al., 1996, 1999). As in the mouse, the number of IAHCs significantly exceeds the low numbers of HSCs in the AGM (Ivanovs et al., 2011), suggesting that many cells in human IAHCs could be nascent HSCs, that cannot yet be detected by direct transplantation, as inferred from *in vitro* modelling of mouse HSC development (Taoudi et al., 2008; Rybtsov et al., 2011; Zhou et al., 2016). The development of a similar *in vitro* co-culture approach for recapitulating human HSC development has been unexpectedly challenging.

Xenotransplantation into immunodeficient mice revealed that the earliest human HSCs emerging in the AGM region are CD34<sup>+</sup>VE-cadherin<sup>+</sup>CD45<sup>+</sup>, consistent with an endothelial origin (North et al., 2002; Taoudi et al., 2005; Ivanovs et al., 2014). This population comprises 500–1000 cells and is uniformly labelled by other UCB HSC markers (Majeti et al., 2007; Notta et al., 2011), which together localise the first few definitive human HSCs to CD34<sup>+</sup>VE-cadherin<sup>+</sup>CD45<sup>+</sup>KIT<sup>+</sup>THY1<sup>+</sup>endoglin<sup>+</sup>RUNX1<sup>+</sup>CD38<sup>−/lo</sup>CD45RA<sup>−</sup> cells (Ivanovs et al., 2014). Thus, compared with UCB from which HSCs can be isolated with high purity (Notta et al., 2011), identification of definitive HSCs within the AGM population is inefficient, and remains a challenge. Phenotypic and quantitative characterisation of the entire HSC hierarchy emerging in the human AGM region is hampered by the lack of an experimental system to identify immature human HSCs. Dissecting the composition of the AGM-derived cell population that contains HSCs using additional – and currently unknown – markers might provide further insights.

Human foetal liver and UCB HSCs could also be a source of useful information for the identification of immature HSCs in the AGM region (Notta et al., 2011; Prasad et al., 2015). In human, the phenotypic identity of foetal liver HSCs is less defined than in the mouse (Kim et al., 2006), although their main markers, CD34<sup>+</sup>CD38<sup>−</sup>, are the same as those of UCB and bone marrow HSCs (Majeti et al., 2007; Doulatov et al., 2012). GPI-80 (VNN2) marks HSCs within the CD34<sup>+</sup>CD38<sup>−/lo</sup>CD90 (THY1)<sup>+</sup> foetal liver population and could potentially be useful for enrichment of developing HSCs from earlier pre-liver stages. Similarly, it would be interesting to establish whether  $\alpha 6$  integrin (also known as CD49f), a marker for UCB HSCs, can be used for this purpose (Notta et al., 2011).

### The signalling landscape within the human AGM niche

HSC development is closely associated with the development of the dorsal aorta and these two developmental programmes share common signalling pathways, which have mainly been investigated using model organisms. In various species, KDR/VEGF signalling plays an important role in the formation of the dorsal aorta (Ciau-Uitz et al., 2016). Groups of KDR<sup>+</sup>CD34<sup>−</sup> cells can be detected at CS 10 (21 dpc) in the trunk mesoderm just before the onset of blood circulation (Cortés et al., 1999). These cells could potentially be considered as candidates for the definitive haemangioblasts described in model organisms and in hPSC cultures (Kennedy et al., 2007; Ciau-Uitz et al., 2016; Slukvin, 2016). By CS 14–15 (30–34 dpc), endothelial cells of the dorsal aorta and IAHCs co-express KDR and its ligand VEGF (Marshall et al., 1999) and upregulate transcription factors that are crucially important for haematopoietic development in IAHCs, such as RUNX1 (also known as AML1), c-myb (MYB), SCL (TAL1), GATA2 and GATA3 (Tavian et al., 1996; Labastie et al., 1998; Marshall et al., 1999; Ivanovs et al., 2014).



Ventral polarisation in IAHC and HSC development in model organisms is at least partly driven by spatial asymmetry of molecular signalling in the AGM region (Wilkinson et al., 2009; Souilhol et al., 2016), suggesting similar mechanisms in human. From CS 13 (28 dpc), BMP4 expression is observed ventrally in a thin subendothelial mesenchymal layer of the dorsal aorta, which transiently expands and subsequently disappears by CS 16 (38 dpc). At this stage, functionally defined HSCs are still present in the AGM region (Ivanovs et al., 2011). BMP4 is a negative regulator of HSC maturation: although it is involved in the formation of the HSC niche, its direct action on HSCs is likely limited by BMP antagonists (Souilhol et al., 2016). Notably, BMP4 can induce expression of tenascin C and fibronectin (Molloy et al., 2008), two extracellular matrix molecules that are also ventrally polarised in the human AGM region (Marshall et al., 1999). Tenascin C promotes haematoendothelial development of hESCs (Uenishi et al., 2014) and its deficiency impairs mouse bone marrow cultures, which can be rescued by fibronectin (Ohta et al., 1998). TGF $\beta$ , another well-known modulator of the extracellular matrix, is expressed in close proximity to IAHCs at the same time as BMP4 expression is downregulated. It has been proposed that TGF $\beta$  could participate in rearrangement of the niche and restriction of HSC expansion prior to foetal liver colonisation (Marshall et al., 2000).

Signalling through KIT is important for HSC maturation and its ligand, SCF (KITL in mouse; KITLG in human), is ventrally polarised in the mouse AGM region (Rybtsov et al., 2014; Souilhol et al., 2016). KIT is expressed in IAHCs and in the first functional HSCs emerging in the human embryo (Labastie et al., 1998; Ivanovs et al., 2014). BMP4 can upregulate KIT in various tissues and, through upregulation of KIT in human aortic endothelium, it may potentially facilitate HSC initiation (Marshall et al., 2007). However, further HSC development may require suppression of BMP4 signalling, as described in the mouse (Souilhol et al., 2016). FLT3, which marks mouse embryonic HSC precursors (Boyer et al., 2011), and its ligand are also expressed in IAHCs and surrounding endothelium, and may be involved in human HSC maturation (Marshall et al., 1999).

### Adhesion and migration

HSC development is associated with cell migration and IAHCs are marked by a range of adhesion molecules that potentially play a role in this process. CD34 and its receptor L-selectin (CD62L; SELL) are involved in cell adhesion processes including lymphocyte-endothelial interactions (Rosen, 2004), and are expressed in human embryonic endothelium, IAHCs and functional HSCs (Tavian et al., 1996; Ivanovs et al., 2014). The first lineage-restricted haematopoietic marker, SPN, appears during hPSC differentiation (Vodyanik et al., 2006) and can mediate adhesion through E-selectin receptor (SELE) (Matsumoto et al., 2005). CD34 and CD43 can cooperatively define adhesive behaviour of haematopoietic cells (Drew et al., 2005). Other surface molecules involved in cell adhesion – ALCAM, VE-cadherin, CD44, CD164 and VCAM1 – are also expressed in IAHCs, endothelial cells of the dorsal aorta and sometimes the mesenchymal cells underneath IAHCs (Cortés et al., 1999; Watt et al., 2000; Marshall, 2006; Ivanovs et al., 2014). Given the role of the TNF/lymphotoxin/NF $\kappa$ B pro-inflammatory pathway in early HSC development described in model organisms (Espín-Palazón et al., 2014; Li et al., 2014) and its impact on VCAM1, L- and E-selectin expression (Suna et al., 2008), it is conceivable that complex and finely regulated adhesion dynamics play important roles in human HSC development. However, it is likely that additional signalling pathways described in other model

organisms (Ciau-Uitz et al., 2016) also play a role in human HSC development. Modelling HSC development *in vitro* using PSCs and other approaches could shed light on this issue.


### Recapitulation of human haematopoietic development using pluripotent stem cells

This section examines parallels between the *in vitro* haematopoietic differentiation of hPSCs and aspects of haematopoietic development in the human embryo. Rather than replicating recent reviews that comprehensively discuss protocols for the differentiation of hPSCs into HSCs or progenitors (Ditadi et al., 2016; Slukvin, 2016; Wahlster and Daley, 2016), we have structured this section to highlight the extent to which *in vitro* differentiation reflects human haematopoietic development (Fig. 4). We argue that ongoing efforts to generate HSCs from PSCs *in vitro* are both informed by and add to our understanding of the signalling events that underpin *in vivo* human haematopoietic development.

### Mesoderm induction and patterning

In the mammalian embryo, ingression of cells through the primitive streak during gastrulation generates mesoderm, which colonises the yolk sac (extra-embryonic mesoderm) and intra-embryonic sites including paraxial, intermediate and lateral plate mesoderm (Tam and Behringer, 1997). Whereas extra-embryonic mesoderm gives rise to transient yolk sac haematopoiesis, lateral plate mesoderm adjacent to the paraxial mesoderm (the splanchnopleura) forms bilateral mesodermal strips that converge to the midline where they coalesce to form the dorsal aorta. The growth factors involved in mesoderm specification, dorsal aorta formation and haematopoietic development include members of the FGF, WNT, TGF $\beta$ , retinoic acid, Hedgehog and Notch signalling pathways (Lawson et al., 2001; Gering and Patient, 2005; Chanda et al., 2013; Lizama et al., 2015; Ciau-Uitz et al., 2016).

In culture, as *in vivo*, mesoderm induction and patterning is mediated by the cooperative actions of highly conserved FGF, BMP4, activin and canonical WNT signals (Schier and Shen, 2000; Kimelman, 2006; Nostro et al., 2008; Woll et al., 2008; Wang and Nakayama, 2009; Bernardo et al., 2011; Yu et al., 2011). Mesoderm patterned to a haematopoietic fate is marked by the expression of primitive streak genes, such as the transcription factor genes *brachyury* (*T*), *MIXL1*, *FOXF1*, as well as cell surface receptors *KDR* and *PDGFRA* (Davis et al., 2008; Ditadi et al., 2016; Slukvin, 2016). An early study showed that the most efficient production of haematopoietic cells resulted from the exposure of the emerging mesoderm to activin and BMP4 signalling (Nostro et al., 2008), analogous to the posterior primitive streak from whence the first yolk sac haematopoietic progenitors are derived. Arguably, this might explain why the haematopoietic output also resembled that of the yolk sac and did not include long-term HSCs. It was only later that a series of studies shed light on the role of mesoderm patterning for both the suppression of primitive haematopoiesis and for the support of definitive haematopoietic lineages. Following an initial period of BMP4-based mesoderm induction, brief inhibition of activin (Kennedy et al., 2012), or provision of a WNT agonist (Gertow et al., 2013) were sufficient to inhibit the erythroid-biased primitive programme. Importantly, activin inhibition (Kennedy et al., 2012), WNT stimulation (Sturgeon et al., 2014) or a combination of the two (Ng et al., 2016) from days 2–4 of *in vitro* differentiation also resulted in a bias towards definitive haematopoietic lineages, as defined by the capacity to generate T lymphocytes (Kennedy et al., 2012; Sturgeon et al., 2014).



**Fig. 4. Model of *in vitro* haematopoietic differentiation from human pluripotent stem cells.** Separation between extra-embryonic, yolk sac-like haematopoiesis (blue shading and arrows) and intra-embryonic AGM-like haematopoiesis (red shading and arrows) initially occurs at the time of mesoderm patterning [days (d) 2-4 of differentiation]. WNT agonists (CHIR99021, denoted CHIR) and activin antagonists (SB431542, denoted SB) orchestrate this switch, although other molecules, such as retinoic acid, may also play a role. A key difference between the extra- and intra-embryonic programmes is the expression of a specific combination of *HOXA* genes that is restricted to the endothelia and haematopoietic progenitors of intra-embryonic haematopoiesis. The extra-embryonic programme generates blast colony-forming cells (BL-CFCs), which are suppressed in the intra-embryonic programme. In both programmes, haemogenic endothelia (HE) generate erythro-myeloid progenitors (EMPs) and T lymphocytes. Thus far, B cells have been generated from an extra-embryonic type protocol although it is anticipated that similar B cells can be derived from AGM-like HE (dashed red arrow). The asterisk indicates that it is not known whether the B cells generated from hPSCs are B-1 or B-2 type. The intra-embryonic AGM-like HE yields cells that have pre-HSC characteristics. Although HSCs and their progeny are predicted (greyed arrows and images), these have yet to be formally shown.

The selective expression of *HOXA* genes in human foetal liver and UCB-derived haematopoietic progenitor and stem cells clearly distinguished them from PSC-derived haematopoietic progenitors generated using conventional mesoderm induction protocols (Dou et al., 2016; Ng et al., 2016). *HOX* genes mark axial position during development and are first expressed at the primitive streak stage (Deschamps and van Nes, 2005). Therefore, it was relevant that activin inhibition and WNT stimulation during mesoderm patterning, alone and in combination, increased the expression of the *CDX* genes, leading to sustained expression of their target *HOXA* genes and more closely mimicking the *HOXA* expression signature seen in foetal liver and cord blood cells (Ng et al., 2016). These data support the premise that allocation of cells to extra- and intra-embryonic haematopoietic fates is initiated during mesoderm patterning, and that this is reflected in the selective expression of *HOXA* genes in definitively patterned mesoderm (Fig. 4).

#### The generation of yolk sac-like haematopoietic lineages

The emergence of FGF2- and VEGF-dependent blast colonies from BMP4-induced mesoderm after 2-4 days of differentiation marks the onset of haematopoiesis in human hPSC cultures (Kennedy et al., 2007; Davis et al., 2008; Vodyanik et al., 2010). As well as the surface markers KDR and PDGFR $\alpha$ , blast colony-forming cells (BL-CFCs) also express the apelin receptor (Vodyanik et al., 2010; Choi et al., 2012; Yu et al., 2012), and CD235a (glycophorin A) (Sturgeon et al., 2014). As in the mouse (Huber et al., 2004), human blast colonies give rise to haematopoietic, endothelial and smooth muscle lineages (Yu et al., 2012). Haematopoietic differentiation of blast colonies is limited to primitive erythrocytes, megakaryocytes and macrophages, and, as in the mouse (Lancrin et al., 2009), probably occurs via a haematogenic endothelial cell precursor.

Further differentiation of hPSCs along a yolk sac-type path can be achieved by co-culture with mouse OP9 stromal cells (Vodyanik et al., 2006), as embryoid bodies (Pick et al., 2007; Nostro et al., 2008) or monolayer cultures (Uenishi et al., 2014) in serum-free media supplemented with growth factors, usually including VEGF, SCF and FGF2. Endothelial and haematopoietic cell surface markers are sequentially acquired, starting with CD34 after 4-6 days, and followed by CD43 at day 6-8, the expression of which first distinguishes haematopoietic from endothelial lineages (Vodyanik et al., 2006). The formation of these blood cells from an endothelial intermediate indicates that the endothelial-to-haematopoietic transition is a common mechanism for blood formation both in the embryo and in differentiating hPSCs and is not limited to a particular developmental stage.

The CD34<sup>+</sup>CD43<sup>+</sup> haematopoietic progenitors give rise to a broad range of erythroid and myeloid lineages (Choi et al., 2012; Kennedy et al., 2012; Ng et al., 2016), reminiscent of the broad potential of the yolk sac-derived erythro-myeloid progenitor (EMP) described in the mouse that transiently seeds the foetal liver (McGrath et al., 2015). Although EMP-type haematopoiesis has not been formally characterised in human embryos, kinetic studies report a rapid transition of clonogenic cells from the yolk sac to the foetal liver at around 5 weeks of gestation (Migliaccio et al., 1986). Given that at this time the human foetal liver does not yet possess repopulating ability (Ivanovs et al., 2011), the cells seeding the early foetal liver are likely to include yolk sac-derived EMPs, and the yolk sac-like erythroid and myeloid cells derived from hPSCs may be their human *in vitro* equivalent. Whether these EMP-type cells are defined as a second 'primitive' or a first 'definitive' haematopoietic wave is somewhat moot. For the moment, the descriptor 'EMP' may be sufficient. We do point out, however, that in contrast to haematopoietic cells derived from the AGM region, foetal liver or



other embryonic sites, hPSC-derived EMP-like cells do not express *HOXA* genes (Dou et al., 2016; Ng et al., 2016).

### The generation of AGM-like haematopoietic lineages from haematogenic endothelium

As discussed above, the appearance of IAHCs in the AGM region, which arise from a transient haematogenic endothelium in the dorsal aorta, predicts the emergence of the first stem cells that acquire repopulating activity in the vertebrate embryo. In contrast to non-haematogenic vascular cells, hPSC-derived haematogenic endothelium lacks expression of CD73 (NT5E) (Choi et al., 2012; Ditadi et al., 2015), CXCR4 and DLL4 (Ditadi et al., 2015). Also, both mouse (North et al., 1999; Thambyrajah et al., 2016) and human (Choi et al., 2012; Ng et al., 2016) haematogenic endothelium are distinguished by their expression of the key transcription factors RUNX1 and GFI1, which are required for the endothelial to haematopoietic transition. Interestingly, hPSCs differentiated with a protocol that induces *HOXA* gene expression produce SOX17-expressing endothelial cells that bear similarity at the transcriptional level to developing dorsal aorta endothelium, and a SOX17<sup>dull</sup> haematogenic endothelial population that efficiently generates haematopoietic cells (Ng et al., 2016). The low level of *SOX17* in haematogenic cells is consistent with literature reporting the downregulation of *SOX17* during the EHT (Clarke et al., 2013; Nobuhisa et al., 2014). Similarly, single-cell analysis of hPSC-derived cells documented a downregulation of *SOX17* at the EHT point (Guibentif et al., 2017).

It has not been possible to identify cell surface markers that distinguish true or 'definitive' AGM-derived HSCs from yolk sac-type, non-engrafting haematopoietic progenitors (Ditadi et al., 2016; Slukvin, 2016). For this reason, considerable attention has been paid to the generation of lymphocytes from hPSCs *in vitro*, arguing that these lineages represent definitive haematopoiesis and may indicate culture conditions that support the development of AGM-like HSCs. Many laboratories have generated T cells from hPSCs, in a Notch-dependent fashion, from endothelial precursors present at 7–9 days of differentiation (Timmermans et al., 2009; Kennedy et al., 2012; Uenishi et al., 2014; Ditadi et al., 2015; Dou et al., 2016; Ng et al., 2016). The dependence of AGM-type, but not yolk sac-type haematopoiesis on Notch signals in the mouse embryo (Kumano et al., 2003; Hadland et al., 2004) has been used to further strengthen the case that the hPSC-derived T cells are progeny of an intra-embryonic haematopoietic progenitor. An important caveat, however, is the realisation that, in the mouse, both T- and B-cell potential is present in the pre-AGM stage yolk sac (Yoshimoto et al., 2011, 2012; McGrath et al., 2015). The dependence of human yolk sac-derived EMP or lymphoid cells on Notch signals is not known.

Unlike the relative ease in which T cells can be derived, B-cell development from hPSCs has been infrequently reported. One laboratory described the generation of cells with a CD19<sup>+</sup>CD10 (MME)<sup>+</sup> pre-B-cell phenotype, some of which matured to surface IgM<sup>+</sup> immature B cells (French et al., 2015).

To place this in context, during mouse development there are several waves of B-cell production (Herzenberg and Herzenberg, 1989). The first innate type B cells, called B-1 cells, arise independently from haematogenic endothelia in both the yolk sac and the AGM region, prior to HSCs, and seed the foetal liver (Yoshimoto, 2015). Subsequent waves of HSC-derived B-1 and adaptive B-2 lymphocytes arise later in the foetal liver (Yoshimoto, 2015) and a wave of predominantly HSC-derived B-2 lymphocytes emerges in the bone marrow (Montecino-Rodriguez and Dorshkind, 2012). The identification of B-1 lymphocytes in humans (Rothstein

and Quach, 2015), and their observation in the 10–11 week foetal liver (Bueno et al., 2016), argues for conservation of the innate immune system between species. Given that human B-1 cells differ immunophenotypically from B-2 lymphocytes (Rothstein and Quach, 2015), it should therefore be possible to discern whether B cells derived from human hPSCs include B-2 cells, and are thus unequivocally descended from HSCs.

The pattern of globin chain expression by hPSC-derived erythroid cells has also served as an indicator of developmental stage. Erythroid cells differentiated from human PSCs usually express  $\epsilon$ - and  $\gamma$ -globins (Chang et al., 2006; Lu et al., 2008; Hatzistavrou et al., 2009; Dias et al., 2011), although prolonged culture in human plasma or on stromal layers has yielded predominantly  $\gamma$ - or  $\gamma$ - and  $\beta$ -globin-expressing cells (Qiu et al., 2008; Lapillonne et al., 2010; Yang et al., 2014; Fujita et al., 2016). Importantly, the switch to  $\beta$ -globin synthesis occurs perinatally, and thus is not expected at the AGM or foetal liver stages of haematopoiesis. Rather, focus is on the anticipated downregulation of embryonic  $\epsilon$ - and  $\zeta$ -globins that signifies exit from primitive haematopoiesis. As a further complication, yolk sac EMP-derived erythroid precursors in mice carrying a single copy of the human  $\beta$ -globin locus downregulate  $\epsilon$ -globin and express low levels of  $\beta$ -globin when matured *in vitro* (McGrath et al., 2011). As with the generation of lymphoid cells, these findings indicate that it is not easy to exclude yolk sac EMP haematopoiesis as a source of erythroid cells that appear to derive from foetal liver because they have suppressed embryonic globins.

Finally, although EMPs are traditionally associated with the yolk sac, cells with a similar restricted developmental capacity could conceivably derive from the AGM haematogenic endothelium also. Consistent with this hypothesis, clonogenic progenitors (approximately 90 colonies per embryo) are present in the VE-cadherin<sup>+</sup>CD45<sup>+</sup> population in the human AGM region, the same fraction harbouring less frequent pre-HSCs and HSCs (Ivanovs et al., 2014). These clonogenic cells might also seed the early foetal liver, and might only be distinguishable from yolk sac-derived EMPs by their expression of *HOXA* genes. Although further experiments will be required as proof, there is support for this idea in a recent report of  $\epsilon$ - to  $\gamma$ -globin switching and downregulation of  $\zeta$ -globin in clonogenic cells derived from *HOXA*<sup>+</sup> cultures (Ng et al., 2016).

Transcriptional comparisons of cells emerging from hPSC-derived *HOXA*<sup>+</sup> haematogenic endothelium and AGM-derived stem progenitors has revealed a high degree of similarity across a broad range of haematopoietic cell surface markers, transcription factors and signalling molecules (Ng et al., 2016). Nevertheless, the hPSC-derived progenitors lack some essential characteristics that enable long-term haematopoietic repopulation. Although mesoderm patterning with modulation of WNT and activin signals increases *HOXA* expression, the pattern of expression in the hPSC-derived progenitors differs subtly from that observed in human AGM samples, with reduced expression of anterior *HOXA* genes and elevated levels of the most posterior *HOXA* genes (Ng et al., 2016).

### Alternative strategies to explore human haematopoietic development

#### Teratoma formation as a model for human HSC development

*In vivo* teratoma formation may offer valuable insights into the generation of transplantable HSCs from hPSCs. Teratomas, which contain derivatives of all three germ layers, can be formed after ectopic engraftment of PSCs into a host animal. The mixture of cell

types in these tumours can preserve some developmentally important inductive interactions, leading to the formation of disorganised tissues and organs, including bone marrow (Hentze et al., 2009; Amabile et al., 2013). Teratomas formed from hPSCs engrafted into immunodeficient mice, in some cases, generate human CD45<sup>+</sup> cells that home to bone marrow, circulate in the recipient peripheral blood and colonise lymphoid tissues (Amabile et al., 2013; Suzuki et al., 2013). Transplantation of total bone marrow or sorted human CD34<sup>+</sup>CD45<sup>+</sup> bone marrow cells from primary into secondary recipients shows donor-derived multilineage peripheral blood chimerism, comparable to UCB repopulation (Amabile et al., 2013; Suzuki et al., 2013). Teratoma-derived B and T cells appear functionally normal, and erythrocytes express adult haemoglobins, which is not usually achievable during *in vitro* hPSC differentiation. It is conceivable that *in vivo* teratoma formation might more faithfully recapitulate some of the processes that occur during embryo development and, in some cases, this could include the specification of an AGM-like region and lead to the production of HSCs. However, an extended period of observation is required to determine whether these HSC-like cells have a truly long-term haematopoietic potential (Ivanovs et al., 2011; Notta et al., 2011).

#### Direct cell lineage conversion: programming and reprogramming to blood


Although enforced gene expression has been shown to be a powerful tool for altering cell fate for several decades (Davis et al., 1987; Kullessa et al., 1995), the real boost to the generation of blood cells using this approach was precipitated by reprogramming studies in which fibroblasts were reverted back to pluripotency using four transcription factors (Takahashi and Yamanaka, 2006; Pereira et al., 2012). These studies spawned quests for combinations of transcription factors, overexpression of which could generate clinically relevant cell types, including HSCs. Groups chose either ESCs (Kyba et al., 2002; Elcheva et al., 2014), human endothelial cells (Sandler et al., 2014), haematopoietic cells (Doulatov et al., 2013) or fibroblasts (Pereira et al., 2013; Batta et al., 2014) as the starting material. Comparative transcriptomics of HSC and progenitor cell populations often provided lists of candidate factors to overexpress. Although many of the factors used in reprogramming studies have previously been implicated in developmental haematopoiesis, their true ‘haematopoietic potency’ becomes unveiled when they are expressed in a foreign cellular context, such as when reprogramming fibroblasts (Pereira et al., 2013; Batta et al., 2014) or undifferentiated PSCs (Elcheva et al., 2014; Vereide et al., 2014) (see Fig. 5 for details).

Using an hPSC differentiation model, the potency of two pairs of transcription factors, ETV2/GATA2 and GATA2/SCL, has been shown in activating pan-myeloid or erythro-megakaryocytic programmes, respectively (Elcheva et al., 2014). These results reflect the importance of these factors as haematopoietic fate-determining transcription factors (Elcheva et al., 2014). Although this system failed to generate long-term repopulating cells, it is a powerful tool for the dissection of haematopoietic developmental pathways. In separate studies, a HOXA9/ERG/RORA/SOX4/MYB combination of transcription factors was identified by comparing a human HSC-enriched population with more mature progenitors from UCB (Doulatov et al., 2010; Laurenti et al., 2013). Overexpression of this combination of transcription factors in hPSC-derived haematopoietic progenitors conferred transient *in vivo* engraftment capacity (Doulatov et al., 2013). The data suggested that HOXA9, ERG, RORA, SOX4 and MYB belong to

the same molecular regulatory network; this is supported by the observation that HOXA9 occupies *Erg*, *Sox4* and *Myb* promoters (Huang et al., 2012). The interest in analysis of HOX genes reflects their strong positive impact on expansion of mouse bone marrow HSCs (Antonchuk et al., 2002; Argiropoulos and Humphries, 2007). Moreover, collinear activation of HOX genes is involved in temporal regionalisation of the mesoderm (Iimura and Pourqu  , 2006), which may underlie functional differences in sequentially emerging haematopoietic waves. For example, HoxB4 overexpression along with adjusted Notch signalling confers multilineage long-term repopulation ability to mouse ESCs (Lu et al., 2016). Downregulation of HOXA5 and HOXA7 in human foetal liver HSCs suppresses their capacity to self-renew and repopulate NSG recipients. Although, forced expression of HOXA5, HOXA7 and HOXA9 alone or in combination in HOXA-negative hPSC-derived progenitors was insufficient to convert the progenitors to HSCs (Dou et al., 2016), a recent study reported a greater degree of success by using a larger cohort of genes (Sugimura et al., 2017). Induced expression of seven transcription factors, notably including three *HOXA* genes, successfully converted hPSC-derived haematogenic endothelium into long-term repopulating HSCs (Sugimura et al., 2017). Finally, expression of four transcription factors (FOSB, GF11, RUNX1 and SPI1) in human umbilical vein and dermal microvascular endothelial cells, or in adult mouse endothelial cells, were also sufficient to generate immunocompetent HSCs (Sandler et al., 2014; Lis et al., 2017). In all these studies, using transcription factors to manipulate cellular identity has provided insights into the regulation of normal haematopoietic development (Fig. 5). For example, successful reprogramming of fibroblasts into blood allowed, through obtaining a molecular signature, identification of precursors of long-term repopulating cells in the mouse placenta (Pereira et al., 2016).

#### Future directions and challenges

Understanding the development of the human haematopoietic system is of fundamental biological and clinical importance. Various haematopoietic diseases originate prenatally (Hunger et al., 1998; Greaves, 2005; Hong et al., 2008; Roy et al., 2012; Barrett et al., 2016), and oncogenic mutations in developing haematopoietic progenitors or stem cells can lead to pre-leukaemic changes that may not be manifested if the oncogene is activated in the adult counterparts (Barrett et al., 2016). The molecular mechanisms that regulate development and maintenance of haematopoietic progenitor and stem cells in the embryo differ significantly from those in the adult, and the functional heterogeneity that appears within the emerging HSC population during development further complicates our understanding of HSC regulation (Copley et al., 2012; Crisan et al., 2015; Beaudin et al., 2016). Although the impetus to study human haematopoietic development is clear, poor availability and heterogeneity of human embryonic material makes the analyses challenging. Direct extrapolation of results obtained from model organisms is not always appropriate owing to species-specific differences, although major genetic networks and sites of haematopoiesis are significantly conserved. Furthermore, for ethical and practical reasons, studying the period of early implantation in the human is extremely difficult. Thus, this highly relevant developmental stage remains largely inaccessible and so cannot inform protocols for derivation of HSCs from PSCs. Additionally, until recently, PSC differentiation was biased towards transitory yolk sac haematopoiesis, resulting in the failure of this *in vitro* model to fully inform us of the mechanisms of HSC development *in vivo*. Thus, deficiency in our *in vivo*



**Fig. 5. Programming and reprogramming towards haematopoietic fate.** *In vitro* generation of haematopoietic cells usually occurs via either directed differentiation of PSCs (green arrows), or via transcription factor-based reprogramming of a variety of different starting cell types (blue boxes). These include ESCs and iPSCs, human umbilical vein endothelial cells (HUVECs), adult fibroblasts, haematogenic endothelium cells, and pro-B cells. Transcription factor combinations used for reprogramming as well as the direction of reprogramming are shown in red (human) and purple (mouse). ETO is also known as RUNX1T1. References are indicated: (1) (Vereide et al., 2014); (2) (Kyba et al., 2002; Lu et al., 2016); (3) (Doulatov et al., 2013); (4) (Elcheva et al., 2014); note that GATA2/SCL combination induces erythro-megakaryocytic differentiation (not shown); (5) (Sandler et al., 2014); (6) (Riddell et al., 2014); (7) (Pereira et al., 2013); (8) (Batta et al., 2014); (9) (Cheng et al., 2016); (10) (Pulecio et al., 2014); (11) (Lis et al., 2017); (12) (Sugimura et al., 2017). CLPs, common lymphoid progenitors; CMPs, common myeloid progenitors; MPPs, multipotent progenitors.

knowledge has not yet been compensated for by information from the *in vitro* system, and vice versa.

In our opinion, two approaches may be taken to address this problem. Notwithstanding their limitations, it is important to continue to use model organisms, which remain virtually unrestricted resources in which to pursue detailed analyses of HSC development. Although not free from ethical and practical considerations, research using non-human primates may also provide valuable information that will help to bridge the evolutionary gap between the mouse and human (D'Souza et al., 2016).

A second approach is the continued use of reprogramming to investigate key determinants of cell fate. Although reprogramming seems like a bold approach to generating cells of interest, it could potentially inform us about events that occur *in vivo*, provided that it goes beyond describing acquisition of markers and extends to functional characteristics. Research in this direction should aim to identify the best cell sources and factors for producing normal, functional HSCs. To date, the selection of genes for haematopoietic reprogramming has been based on genes known to be essential for haematopoietic development and genes that are differentially expressed in bone marrow/foetal liver/UCB HSCs compared with progenitor cells. However, the genes that drive the formation of the first exceptionally potent HSCs that emerge in the human AGM region (Ivanovs et al., 2011) remain unknown. Identification of

these genes in human requires more detailed reconstruction of the developing HSC hierarchy, which is currently limited by lack of an appropriate experimental system in human, but could be facilitated by further in-depth analysis of model organisms. Although the data show a lack of definitive HSCs in human placenta and large extra-embryonic arteries, they should not be excluded from consideration as they could harbour immature HSCs.

The presence of one or more transgenes in the genome of reprogrammed human HSCs represents a clinical risk, and further makes it difficult to assess the entire scale of molecular changes in the reprogrammed cells. Reprogramming methods that do not involve transgene integration, such as non-integrating vectors and RNA, will therefore be highly desirable. Whether this approach, which has proved suitable for iPSC generation, will work for reprogramming into HSCs will depend on the activation of crucial endogenous genes that act as 'molecular switches' to produce a phenotypically stable cell type, long after the non-integrating factors are degraded or diluted. This is particularly important for conferring stemness to progenitor cells, but so far it is not clear whether this approach will be successful. Ultimately, the safest HSCs generated in the dish would likely be genetically unaltered cells, produced by exposure to external factors equivalent to the embryonic cues that drive HSC development. Here, the bottleneck seems to be identifying the combinations of soluble and cell-based factors and



inhibitors that will produce bona fide long-term repopulating human HSCs and again, analysis of mouse and other model organisms as well as reprogramming studies may shed light on this problem. The generation of reporter hPSC lines will continue to be a powerful methodology for analysing human HSC development (Raffi et al., 2013; Jung et al., 2016; Ng et al., 2016). Ultimately, the depth of our understanding of human HSC development will be measured by our ability to faithfully recapitulate *in vitro* the embryonic pathway that leads to HSC formation. This is what will pave the way to informative *in vitro* modelling of childhood haematological pathologies, and to the robust generation of bona fide HSCs for possible clinical use.

#### Acknowledgements

The authors would like to thank Prof. I. Slukvin (Madison) for helpful comments on hPSC differentiation; Dr D. Paruzina for providing images for Fig. 2C; Dr S. Gordon-Keylock for helpful comments and assistance during preparation of this manuscript.

#### Competing interests

The authors declare no competing or financial interests.

#### Funding

This work was supported by Bloodwise (formerly LLR) and Medical Research Council (UK). E.G.S. and A.G.E. are Research Fellows of the National Health and Medical Research Council (Australia). Work in the laboratories of E.G.S. and A.G.E. is supported by the National Health and Medical Research Council (Australia), Stem Cells Australia and the Children's Cancer Foundation.

#### References

- Amabile, G., Welner, R. S., Nombela-Arrieta, C., D'Alise, A. M., Di Ruscio, A., Ebralidze, A. K., Kraytsberg, Y., Ye, M., Kocher, O., Neuberg, D. S. et al. (2013). In vivo generation of transplantable human hematopoietic cells from induced pluripotent stem cells. *Blood* **121**, 1255–1264.
- Antonchuk, J., Sauvageau, G. and Humphries, R. K. (2002). HOXB4-induced expansion of adult hematopoietic stem cells *ex vivo*. *Cell* **109**, 39–45.
- Argiropoulos, B. and Humphries, R. K. (2007). Hox genes in hematopoiesis and leukemogenesis. *Oncogene* **26**, 6766–6776.
- Barcena, A., Muench, M. O., Kapidzic, M. and Fisher, S. J. (2009). A new role for the human placenta as a hematopoietic site throughout gestation. *Reprod. Sci.* **16**, 178–187.
- Barrett, N. A., Malouf, C., Kapeni, C., Bacon, W. A., Giotopoulos, G., Jacobsen, S. E. W., Huntly, B. J. and Ottersbach, K. (2016). Mli-AF4 confers enhanced self-renewal and lymphoid potential during a restricted window in development. *Cell Rep.* **16**, 1039–1054.
- Batta, K., Florkowska, M., Kouskoff, V. and Lacaud, G. (2014). Direct reprogramming of murine fibroblasts to hematopoietic progenitor cells. *Cell Rep.* **9**, 1871–1884.
- Beaudin, A. E., Boyer, S. W., Perez-Cunningham, J., Hernandez, G. E., Derderian, S. C., Juijavarapu, C., Aaserude, E., MacKenzie, T. and Forsberg, E. C. (2016). A transient developmental hematopoietic stem cell gives rise to innate-like B and T cells. *Cell Stem Cell* **19**, 768–783.
- Bernardo, A. S., Faial, T., Gardner, L., Nakan, K. K., Ortmann, D., Senner, C. E., Callery, E. M., Trotter, M. W., Hemberger, M., Smith, J. C. et al. (2011). BRACHYURY and CDX2 mediate BMP-induced differentiation of human and mouse pluripotent stem cells into embryonic and extraembryonic lineages. *Cell Stem Cell* **9**, 144–155.
- Bertrand, J. Y., Giroux, S., Golub, R., Klaine, M., Jalil, A., Boucontet, L., Godin, I. and Cumano, A. (2005). Characterization of purified intraembryonic hematopoietic stem cells as a tool to define their site of origin. *Proc. Natl. Acad. Sci. USA* **102**, 134–139.
- Bertrand, J. Y., Chi, N. C., Santoso, B., Teng, S., Stainier, D. Y. R. and Traver, D. (2010). Haematopoietic stem cells derive directly from aortic endothelium during development. *Nature* **464**, 108–111.
- Bloom, B. and Bartelmez, G. W. (1940). Hematopoiesis in young human embryos. *Am. J. Anat.* **67**, 21–53.
- Böiers, C., Carrelha, J., Lutteropp, M., Luc, S., Green, J. C. A., Azzoni, E., Woll, P. S., Mead, A. J., Hultquist, A., Swiers, G. et al. (2013). Lymphomyeloid contribution of an immune-restricted progenitor emerging prior to definitive hematopoietic stem cells. *Cell Stem Cell* **13**, 535–548.
- Boyer, S. W., Schroeder, A. V., Smith-Berdan, S. and Forsberg, E. C. (2011). All hematopoietic cells develop from hematopoietic stem cells through Flk2/Flt3-positive progenitor cells. *Cell Stem Cell* **9**, 64–73.
- Bueno, C., van Roon, E. H. J., Muñoz-López, A., Sanjuan-Pla, A., Juan, M., Navarro, A., Stam, R. W. and Menendez, P. (2016). Immunophenotypic analysis and quantification of B-1 and B-2 B cells during human fetal hematopoietic development. *Leukemia* **30**, 1603–1606.
- Catlin, S. N., Busque, L., Gale, R. E., Gutter, P. and Abkowitz, J. L. (2011). The replication rate of human hematopoietic stem cells *in vivo*. *Blood* **117**, 4460–4466.
- Chanda, B., Ditadi, A., Iscove, N. N. and Keller, G. (2013). Retinoic acid signaling is essential for embryonic hematopoietic stem cell development. *Cell* **155**, 215–227.
- Chang, K.-H., Nelson, A. M., Cao, H., Wang, L., Nakamoto, B., Ware, C. B. and Papayannopoulou, T. (2006). Definitive-like erythroid cells derived from human embryonic stem cells coexpress high levels of embryonic and fetal globins with little or no adult globin. *Blood* **108**, 1515–1523.
- Charbord, P., Tavian, M., Humeau, L. and Peault, B. (1996). Early ontogeny of the human marrow from long bones: an immunohistochemical study of hematopoiesis and its microenvironment. *Blood* **87**, 4109–4119.
- Chen, M. J., Yokomizo, T., Zeigler, B. M., Dzierzak, E. and Speck, N. A. (2009). Runx1 is required for the endothelial to hematopoietic cell transition but not thereafter. *Nature* **457**, 887–891.
- Chen, M. J., Li, Y., De Obaldia, M. E., Yang, Q., Yzaguirre, A. D., Yamada-Inagawa, T., Vink, C. S., Bhandoola, A., Dzierzak, E. and Speck, N. A. (2011). Erythroid/myeloid progenitors and hematopoietic stem cells originate from distinct populations of endothelial cells. *Cell Stem Cell* **9**, 541–552.
- Cheng, H., Ang, H. Y.-K., El Farran, C. A., Li, P., Fang, H. T., Liu, T. M., Kong, S. L., Chin, M. L., Ling, W. Y., Lim, E. K. H. et al. (2016). Reprogramming mouse fibroblasts into engraftable myeloerythroid and lymphoid progenitors. *Nat. Commun.* **7**, 13396.
- Choi, K.-D., Vodyanik, M. A., Togarrati, P. P., Suknuntha, K., Kumar, A., Samarjeet, F., Probasco, M. D., Tian, S., Stewart, R., Thomson, J. A. et al. (2012). Identification of the hemogenic endothelial progenitor and its direct precursor in human pluripotent stem cell differentiation cultures. *Cell Rep.* **2**, 553–567.
- Ciau-Uitz, A., Patient, R. and Medvinsky, A. (2016). Ontogeny of the hematopoietic system. In *Encyclopedia of Immunobiology*, Vol. 1 (ed. M. J. H. Ratcliffe), pp. 1–14. Oxford: Academic Press.
- Clarke, R. L., Yzaguirre, A. D., Yashiro-Ohtani, Y., Bondue, A., Blanpain, C., Pear, W. S., Speck, N. A. and Keller, G. (2013). The expression of Sox17 identifies and regulates haemogenic endothelium. *Nat. Cell Biol.* **15**, 502–510.
- Copley, M. R., Beer, P. A. and Eaves, C. J. (2012). Hematopoietic stem cell heterogeneity takes center stage. *Cell Stem Cell* **10**, 690–697.
- Cortés, F., Debacker, C., Péault, B. and Labastie, M.-C. (1999). Differential expression of KDR/VEGFR-2 and CD34 during mesoderm development of the early human embryo. *Mech. Dev.* **83**, 161–164.
- Crisan, M. and Dzierzak, E. (2016). The many faces of hematopoietic stem cell heterogeneity. *Development* **143**, 4571–4581.
- Crisan, M., Kartalaei, P. S., Vink, C. S., Yamada-Inagawa, T., Bollerot, K., van Icken, W., van der Linden, R., de Sousa Lopes, S. M. C., Monteiro, R., Mummery, C. et al. (2015). BMP signalling differentially regulates distinct hematopoietic stem cell types. *Nat. Commun.* **6**, 8040.
- Cumano, A., Dieterlen-Lievre, F. and Godin, I. (1996). Lymphoid potential, probed before circulation in mouse, is restricted to caudal intraembryonic splanchnopleura. *Cell* **86**, 907–916.
- Davis, R. L., Weintraub, H. and Lassar, A. B. (1987). Expression of a single transcribed cDNA converts fibroblasts to myoblasts. *Cell* **51**, 987–1000.
- Davis, R. P., Ng, E. S., Costa, M., Mossman, A. K., Sourris, K., Elefanti, A. G. and Stanley, E. G. (2008). Targeting a GFP reporter gene to the MIXL1 locus of human embryonic stem cells identifies human primitive streak-like cells and enables isolation of primitive hematopoietic precursors. *Blood* **111**, 1876–1884.
- de Bruijn, M. F. T. R., Speck, N. A., Peeters, M. C. E. and Dzierzak, E. (2000). Definitive hematopoietic stem cells first develop within the major arterial regions of the mouse embryo. *EMBO J.* **19**, 2465–2474.
- de Bruijn, M. F. T. R., Ma, X., Robin, C., Ottersbach, K., Sanchez, M.-J. and Dzierzak, E. (2002). Hematopoietic stem cells localize to the endothelial cell layer in the midgestation mouse aorta. *Immunity* **16**, 673–683.
- Deschamps, J. and van Nes, J. (2005). Developmental regulation of the Hox genes during axial morphogenesis in the mouse. *Development* **132**, 2931–2942.
- Dias, J., Gumenyuk, M., Kang, H. J., Vodyanik, M., Yu, J., Thomson, J. A. and Slukvin, I. I. (2011). Generation of red blood cells from human induced pluripotent stem cells. *Stem Cells Dev.* **20**, 1639–1647.
- Ditadi, A., Sturgeon, C. M., Tober, J., Awong, G., Kennedy, M., Yzaguirre, A. D., Azzola, L., Ng, E. S., Stanley, E. G., French, D. L. et al. (2015). Human definitive haemogenic endothelium and arterial vascular endothelium represent distinct lineages. *Nat. Cell Biol.* **17**, 580–591.
- Ditadi, A., Sturgeon, C. M. and Keller, G. (2016). A view of human hematopoietic development from the Petri dish. *Nat. Rev. Mol. Cell Biol.* **18**, 56–67.
- Dou, D. R., Calvanese, V., Sierra, M. I., Nguyen, A. T., Minasian, A., Saarikoski, P., Sasidharan, R., Ramirez, C. M., Zack, J. A., Crooks, G. M. et al. (2016). Medial HOXA genes demarcate hematopoietic stem cell fate during human development. *Nat. Cell Biol.* **18**, 595–606.
- Doulatov, S., Notta, F., Eppert, K., Nguyen, L. T., Ohashi, P. S. and Dick, J. E. (2010). Revised map of the human progenitor hierarchy shows the origin of

- macrophages and dendritic cells in early lymphoid development. *Nat. Immunol.* **11**, 585-593.
- Doulatov, S., Notta, F., Laurenti, E. and Dick, J. E. (2012). Hematopoiesis: a human perspective. *Cell Stem Cell* **10**, 120-136.
- Doulatov, S., Vo, L. T., Chou, S. S., Kim, P. G., Arora, N., Li, H., Hadland, B. K., Bernstein, I. D., Collins, J. J., Zon, L. I. et al. (2013). Induction of multipotential hematopoietic progenitors from human pluripotent stem cells via respecification of lineage-restricted precursors. *Cell Stem Cell* **13**, 459-470.
- Drew, E., Merzaban, J. S., Seo, W., Ziltener, H. J. and McNaghy, K. M. (2005). CD34 and CD43 inhibit mast cell adhesion and are required for optimal mast cell reconstitution. *Immunity* **22**, 43-57.
- D'Souza, S. S., Maufort, J., Kumar, A., Zhang, J., Smuga-Otto, K., Thomson, J. A. and Slukvin, I. I. (2016). GSK3beta inhibition promotes efficient myeloid and lymphoid hematopoiesis from non-human primate-induced pluripotent stem cells. *Stem Cell Rep.* **6**, 243-256.
- Eiken, H. M., Nishikawa, S.-I. and Schroeder, T. (2009). Continuous single-cell imaging of blood generation from haemogenic endothelium. *Nature* **457**, 896-900.
- Elcheva, I., Brok-Volchanskaya, V., Kumar, A., Liu, P., Lee, J.-H., Tong, L., Vodnyanik, M., Swanson, S., Stewart, R., Kyba, M. et al. (2014). Direct induction of haematoendothelial programs in human pluripotent stem cells by transcriptional regulators. *Nat. Commun.* **5**, 4372.
- Espín-Palazón, R., Stachura, D. L., Campbell, C. A., García-Moreno, D., Del Cid, N., Kim, A. D., Candel, S., Meseguer, J., Mulero, V. and Traver, D. (2014). Proinflammatory signaling regulates hematopoietic stem cell emergence. *Cell* **159**, 1070-1085.
- Ferkowicz, M. J., Starr, M., Xie, X., Li, W., Johnson, S. A., Shelley, W. C., Morrison, P. R. and Yoder, M. C. (2003). CD41 expression defines the onset of primitive and definitive hematopoiesis in the murine embryo. *Development* **130**, 4393-4403.
- French, A., Yang, C.-T., Taylor, S., Watt, S. M. and Carpenter, L. (2015). Human induced pluripotent stem cell-derived B lymphocytes express sIgM and can be generated via a hemogenic endothelium intermediate. *Stem Cells Dev.* **24**, 1082-1095.
- Fujita, A., Uchida, N., Haro-Mora, J. J., Winkler, T. and Tisdale, J. (2016).  $\beta$ -Globin-expressing definitive erythroid progenitor cells generated from embryonic and induced pluripotent stem cell-derived sacs. *Stem Cells* **34**, 1541-1552.
- Fukuda, T. (1973). Fetal hemopoiesis. I. Electron microscopic studies on human yolk sac hemopoiesis. *Virchows Arch. B Cell Pathol.* **14**, 197-213.
- García-Porrero, J. A., Godin, I. E. and Dieterlen-Lievre, F. (1995). Potential intraembryonic hemogenic sites at pre-liver stages in the mouse. *Anat. Embryol.* **192**, 425-435.
- Gekas, C., Dieterlen-Lievre, F., Orkin, S. H. and Mikkola, H. K. A. (2005). The placenta is a niche for hematopoietic stem cells. *Dev. Cell* **8**, 365-375.
- Gering, M. and Patient, R. (2005). Hedgehog signaling is required for adult blood stem cell formation in zebrafish embryos. *Dev. Cell* **8**, 389-400.
- Gertow, K., Hirst, C. E., Yu, Q. C., Ng, E. S., Pereira, L. A., Davis, R. P., Stanley, E. G. and Elefanty, A. G. (2013). WNT3A promotes hematopoietic or mesenchymal differentiation from hESCs depending on the time of exposure. *Stem Cell Rep.* **1**, 53-65.
- Ghiara, G., Ferkowicz, M. J., Milsom, M. D., Bailey, J., Witte, D., Cancelas, J. A., Yoder, M. C. and Williams, D. A. (2008). Rac1 is essential for intraembryonic hematopoiesis and for the initial seeding of fetal liver with definitive hematopoietic progenitor cells. *Blood* **111**, 3313-3321.
- Greaves, M. (2005). In utero origins of childhood leukaemia. *Early Hum. Dev.* **81**, 123-129.
- Guibentif, C., Rönn, R. E., Böiers, C., Lang, S., Saxena, S., Soneji, S., Enver, T., Karlsson, G. and Woods, N.-B. (2017). Single-cell analysis identifies distinct stages of human endothelial-to-hematopoietic transition. *Cell Rep.* **19**, 10-19.
- Hadland, B. K., Huppert, S. S., Kanungo, J., Xue, Y., Jiang, R., Gridley, T., Conlon, R. A., Cheng, A. M., Kopan, R. and Longmore, G. D. (2004). A requirement for Notch1 distinguishes 2 phases of definitive hematopoiesis during development. *Blood* **104**, 3097-3105.
- Hatzistavrou, T., Micallef, S. J., Ng, E. S., Vadolas, J., Stanley, E. G. and Elefanty, A. G. (2009). ErythRED, a hESC line enabling identification of erythroid cells. *Nat. Methods* **6**, 659-662.
- Hentze, H., Soong, P. L., Wang, S. T., Phillips, B. W., Putti, T. C. and Dunn, N. R. (2009). Teratoma formation by human embryonic stem cells: evaluation of essential parameters for future safety studies. *Stem Cell Res.* **2**, 198-210.
- Herzenberg, L. A. and Herzenberg, L. A. (1989). Toward a layered immune system. *Cell* **59**, 953-954.
- Holyoake, T. L., Nicolini, F. E. and Eaves, C. J. (1999). Functional differences between transplantable human hematopoietic stem cells from fetal liver, cord blood, and adult marrow. *Exp. Hematol.* **27**, 1418-1427.
- Hong, D., Gupta, R., Ancliff, P., Atzberger, A., Brown, J., Soneji, S., Green, J., Colman, S., Piacibello, W., Buckle, V. et al. (2008). Initiating and cancer-propagating cells in TEL-AML1-associated childhood leukemia. *Science* **319**, 336-339.
- Huang, Y., Sitwala, K., Bronstein, J., Sanders, D., Dandekar, M., Collins, C., Robertson, G., MacDonald, J., Cezard, T., Bilensky, M. et al. (2012). Identification and characterization of Hoxa9 binding sites in hematopoietic cells. *Blood* **119**, 388-398.
- Huber, T. L., Kouskoff, V., Fehling, H. J., Palis, J. and Keller, G. (2004). Haemangioblast commitment is initiated in the primitive streak of the mouse embryo. *Nature* **432**, 625-630.
- Hunger, S. P., McGavran, L., Meltesen, L., Parker, N. B., Kassenbrock, C. K. and Bitter, M. A. (1998). Oncogenesis in utero: fetal death due to acute myelogenous leukaemia with an MLL translocation. *Br. J. Haematol.* **103**, 539-542.
- Iimura, T. and Pourquié, O. (2006). Collinear activation of Hoxb genes during gastrulation is linked to mesoderm cell ingression. *Nature* **442**, 568-571.
- Ivanovs, A., Rytsov, S., Welch, L., Anderson, R. A., Turner, M. L. and Medvinsky, A. (2011). Highly potent human hematopoietic stem cells first emerge in the intraembryonic aorta-gonad-mesonephros region. *J. Exp. Med.* **208**, 2417-2427.
- Ivanovs, A., Rytsov, S., Anderson, R. A., Turner, M. L. and Medvinsky, A. (2013). Postmenstrual gestational age should be used with care in studies of early human hematopoietic development. *Blood* **121**, 3051-3052.
- Ivanovs, A., Rytsov, S., Anderson, R. A., Turner, M. L. and Medvinsky, A. (2014). Identification of the niche and phenotype of the first human hematopoietic stem cells. *Stem Cell Rep.* **2**, 449-456.
- Jaffredo, T., Gautier, R., Eichmann, A. and Dieterlen-Lievre, F. (1998). Intraaortic hemopoietic cells are derived from endothelial cells during ontogeny. *Development* **125**, 4575-4583.
- Jaffredo, T., Gautier, R., Brajeul, V. and Dieterlen-Lievre, F. (2000). Tracing the progeny of the aortic hemangioblast in the avian embryo. *Dev. Biol.* **224**, 204-214.
- Jokubaitis, V. J., Sinka, L., Driessen, R., Whitty, G., Haylock, D. N., Bertoncello, I., Smith, I., Peault, B., Tavian, M. and Simmons, P. J. (2008). Angiotensin-converting enzyme (CD143) marks hematopoietic stem cells in human embryonic, fetal, and adult hematopoietic tissues. *Blood* **111**, 4055-4063.
- Jung, H. S., Uenishi, G., Kumar, A., Park, M. A., Raymond, M., Fink, D., McLeod, E. and Slukvin, I. (2016). A human VE-cadherin-tdTomato and CD43-green fluorescent protein dual reporter cell line for study endothelial to hematopoietic transition. *Stem Cell Res.* **17**, 401-405.
- Kauts, M.-L., Vink, C. S. and Dzierzak, E. (2016). Hematopoietic (stem) cell development-how divergent are the roads taken? *FEBS Lett.* **590**, 3975-3986.
- Kennedy, M., D'Souza, S. L., Lynch-Kattman, M., Schwantz, S. and Keller, G. (2007). Development of the hemangioblast defines the onset of hematopoiesis in human ES cell differentiation cultures. *Blood* **109**, 2679-2687.
- Kennedy, M., Awong, G., Sturgeon, C. M., Ditadi, A., LaMotte-Mohs, R., Zúñiga-Pflücker, J. C. and Keller, G. (2012). T lymphocyte potential marks the emergence of definitive hematopoietic progenitors in human pluripotent stem cell differentiation cultures. *Cell Rep.* **2**, 1722-1735.
- Kim, I., He, S., Yilmaz, O. H., Kiel, M. J. and Morrison, S. J. (2006). Enhanced purification of fetal liver hematopoietic stem cells using SLAM family receptors. *Blood* **108**, 737-744.
- Kim, A. D., Stachura, D. L. and Traver, D. (2014). Cell signaling pathways involved in hematopoietic stem cell specification. *Exp. Cell Res.* **329**, 227-233.
- Kimelman, D. (2006). Mesoderm induction: from caps to chips. *Nat. Rev. Genet.* **7**, 360-372.
- Kissa, K. and Herbomel, P. (2010). Blood stem cells emerge from aortic endothelium by a novel type of cell transition. *Nature* **464**, 112-115.
- Kulesha, H., Frampton, J. and Graf, T. (1995). GATA-1 reprograms avian myelomonocytic cell lines into eosinophils, thrombocytes, and erythroblasts. *Genes Dev.* **9**, 1250-1262.
- Kumano, K., Chiba, S., Kunisato, A., Sata, M., Saito, T., Nakagami-Yamaguchi, E., Yamaguchi, T., Masuda, S., Shimizu, K., Takahashi, T. et al. (2003). Notch1 but not Notch2 is essential for generating hematopoietic stem cells from endothelial cells. *Immunity* **18**, 699-711.
- Kumaravelu, P., Hook, L., Morrison, A. M., Ure, J., Zhao, S., Zuyev, S., Ansell, J. and Medvinsky, A. (2002). Quantitative developmental anatomy of definitive haematopoietic stem cells/long-term repopulating units (HSC/RUs), role of the aorta-gonad-mesonephros (AGM) region and the yolk sac in colonisation of the mouse embryonic liver. *Development* **129**, 4891-4899.
- Kyba, M., Perlingeiro, R. C. R. and Daley, G. Q. (2002). HoxB4 confers definitive lymphoid-myeloid engraftment potential on embryonic stem cell and yolk sac hematopoietic progenitors. *Cell* **109**, 29-37.
- Labastie, M. C., Cortes, F., Romeo, P. H., Dulac, C. and Peault, B. (1998). Molecular identity of hematopoietic precursor cells emerging in the human embryo. *Blood* **92**, 3624-3635.
- Lancrin, C., Sroczyńska, P., Stephenson, C., Allen, T., Kouskoff, V. and Lacaud, G. (2009). The haemangioblast generates haematopoietic cells through a haemogenic endothelium stage. *Nature* **457**, 892-895.
- Lapillonne, H., Kobari, L., Mazurier, C., Tropel, P., Giarratana, M.-C., Zanella-Cleon, I., Kiger, L., Wattenhofer-Donze, M., Puccio, H., Hebert, N. et al. (2010). Red blood cell generation from human induced pluripotent stem cells: perspectives for transfusion medicine. *Haematologica* **95**, 1651-1659.
- Laurenti, E., Doulatov, S., Zandi, S., Plumb, I., Chen, J., April, C., Fan, J.-B. and Dick, J. E. (2013). The transcriptional architecture of early human hematopoiesis identifies multilevel control of lymphoid commitment. *Nat. Immunol.* **14**, 756-763.



- Lawson, N. D., Scheer, N., Pham, V. N., Kim, C. H., Chitnis, A. B., Campos-Ortega, J. A. and Weinstein, B. M. (2001). Notch signaling is required for arterial-venous differentiation during embryonic vascular development. *Development* **128**, 3675-3683.
- Li, Y., Esain, V., Teng, L., Xu, J., Kwan, W., Frost, I. M., Yzaguirre, A. D., Cai, X., Cortes, M., Majenburger, M. W. et al. (2014). Inflammatory signaling regulates embryonic hematopoietic stem and progenitor cell production. *Genes Dev.* **28**, 2597-2612.
- Lis, R., Karrasch, C. C., Poulos, M. G., Kunar, B., Redmond, D., Duran, J. G. B., Badwe, C. R., Schachterle, W., Ginsberg, M., Xiang, J. et al. (2017). Conversion of adult endothelium to immunocompetent haematopoietic stem cells. *Nature* **545**, 439-445.
- Lizama, C. O., Hawkins, J. S., Schmitt, C. E., Bos, F. L., Zape, J. P., Cautivo, K. M., Borges Pinto, H., Rhyner, A. M., Yu, H., Donohoe, M. E. et al. (2015). Repression of arterial genes in hemogenic endothelium is sufficient for haematopoietic fate acquisition. *Nat. Commun.* **6**, 7739.
- Lu, S.-J., Feng, Q., Park, J. S., Vida, L., Lee, B.-S., Strausbauch, M., Wettstein, P. J., Honig, G. R. and Lanza, R. (2008). Biologic properties and enucleation of red blood cells from human embryonic stem cells. *Blood* **112**, 4475-4484.
- Lu, Y.-F., Cahan, P., Ross, S., Sahalie, J., Sousa, P. M., Hadland, B. K., Cai, W., Serrao, E., Engelman, A. N., Bernstein, I. D. et al. (2016). Engineered murine HSCs reconstitute multi-lineage hematopoiesis and adaptive immunity. *Cell Rep.* **17**, 3178-3192.
- Luckett, W. P. (1978). Origin and differentiation of the yolk sac and extraembryonic mesoderm in presomite human and rhesus monkey embryos. *Am. J. Anat.* **152**, 59-97.
- Majeti, R., Park, C. Y. and Weissman, I. L. (2007). Identification of a hierarchy of multipotent hematopoietic progenitors in human cord blood. *Cell Stem Cell* **1**, 635-645.
- Marshall, C. (2006). Intraembryonic development of hematopoietic stem cells during human ontogeny. In *Hematopoietic Stem Cell Development* (ed. A. Cumano and I. Godin), pp. 142-153. Berlin, Germany: Springer.
- Marshall, C. J., Moore, R. L., Thorogood, P., Brickell, P. M., Kinnon, C. and Thrasher, A. J. (1999). Detailed characterization of the human aorta-gonad-mesonephros region reveals morphological polarity resembling a hematopoietic stromal layer. *Dev. Dyn.* **215**, 139-147.
- Marshall, C. J., Kinnon, C. and Thrasher, A. J. (2000). Polarized expression of bone morphogenetic protein-4 in the human aorta-gonad-mesonephros region. *Blood* **96**, 1591-1593.
- Marshall, C. J., Sinclair, J. C., Thrasher, A. J. and Kinnon, C. (2007). Bone morphogenetic protein 4 modulates c-Kit expression and differentiation potential in murine embryonic aorta-gonad-mesonephros haematopoiesis in vitro. *Br. J. Haematol.* **139**, 321-330.
- Matsumoto, M., Atarashi, K., Umamoto, E., Furukawa, Y., Shigeta, A., Miyasaka, M. and Hirata, T. (2005). CD43 functions as a ligand for E-Selectin on activated T cells. *J. Immunol.* **175**, 8042-8050.
- McCune, J. M., Namikawa, R., Kaneshima, H., Shultz, L. D., Lieberman, M. and Weissman, I. L. (1988). The SCID-hu mouse: murine model for the analysis of human hematolymphoid differentiation and function. *Science* **241**, 1632-1639.
- McGrath, K. E., Frame, J. M., Fromm, G. J., Koniski, A. D., Kingsley, P. D., Little, J., Bulger, M. and Palis, J. (2011). A transient definitive erythroid lineage with unique regulation of the beta-globin locus in the mammalian embryo. *Blood* **117**, 4600-4608.
- McGrath, K. E., Frame, J. M., Fegan, K. H., Bowen, J. R., Conway, S. J., Catherman, S. C., Kingsley, P. D., Koniski, A. D. and Palis, J. (2015). Distinct sources of hematopoietic progenitors emerge before HSCs and provide functional blood cells in the mammalian embryo. *Cell Rep.* **11**, 1892-1904.
- Medvinsky, A. and Dzierzak, E. (1996). Definitive hematopoiesis is autonomously initiated by the AGM region. *Cell* **86**, 897-906.
- Medvinsky, A. L., Samoylina, N. L., Müller, A. M. and Dzierzak, E. A. (1993). An early pre-liver intraembryonic source of CFU-S in the developing mouse. *Nature* **364**, 64-67.
- Medvinsky, A., Rytsov, S. and Taoudi, S. (2011). Embryonic origin of the adult hematopoietic system: advances and questions. *Development* **138**, 1017-1031.
- Migliaccio, G., Migliaccio, A. R., Petti, S., Mavilio, F., Russo, G., Lazzaro, D., Testa, U., Marinucci, M. and Peschle, C. (1986). Human embryonic hemopoiesis. Kinetics of progenitors and precursors underlying the yolk sac—liver transition. *J. Clin. Invest.* **78**, 51-60.
- Mikkola, H. K. A., Fujiwara, Y., Schlaeger, T. M., Traver, D. and Orkin, S. H. (2003). Expression of CD41 marks the initiation of definitive hematopoiesis in the mouse embryo. *Blood* **101**, 508-516.
- Minot, C. S. (1912). The origin of the angioblast and the development of the blood. In *Manual of Human Embryology*, Vol. 2 (ed. F. Keibel and F. P. Mall), pp. 498-534. Philadelphia: J. B. Lippincott Company.
- Molloy, E. L., Adams, A., Moore, J. B., Masterson, J. C., Madrigal-Estebas, L., Mahon, B. P. and O'Dea, S. (2008). BMP4 induces an epithelial-mesenchymal transition-like response in adult airway epithelial cells. *Growth Factors* **26**, 12-22.
- Montecino-Rodriguez, E. and Dorshkind, K. (2012). B-1 B cell development in the fetus and adult. *Immunity* **36**, 13-21.
- Muench, M. O., Kapidzic, M., Gormley, M., Gutierrez, A. G., Ponder, K. L., Fomin, M. E., Beyer, A. I., Stolp, H., Qi, Z., Fisher, S. J. et al. (2017). The human chorion contains definitive hematopoietic stem cells from the 15th week of gestation. *Development* **144**, 1399-1411.
- Müller, A. M., Medvinsky, A., Strouboulis, J., Grosfeld, F. and Dzierzak, E. (1994). Development of hematopoietic stem cell activity in the mouse embryo. *Immunity* **1**, 291-301.
- Ng, E. S., Azzola, L., Bruveris, F. F., Calvanese, V., Phipson, B., Vlahos, K., Hirst, C., Jokubaitis, V. J., Yu, Q. C., Maksimovic, J. et al. (2016). Differentiation of human embryonic stem cells to HOXA<sup>+</sup> hemogenic vasculature that resembles the aorta-gonad-mesonephros. *Nat. Biotechnol.* **34**, 1168-1179.
- Nobuhisa, I., Osawa, M., Uemura, M., Kishikawa, Y., Anani, M., Harada, K., Takagi, H., Saito, K., Kanai-Azuma, M., Kanai, Y. et al. (2014). Sox17-mediated maintenance of fetal intra-aortic hematopoietic cell clusters. *Mol. Cell. Biol.* **34**, 1976-1990.
- North, T., Gu, T. L., Stacy, T., Wang, Q., Howard, L., Binder, M., Marin-Padilla, M. and Speck, N. A. (1999). Cbfa2 is required for the formation of intra-aortic hematopoietic clusters. *Development* **126**, 2563-2575.
- North, T. E., de Bruijn, M. F. T. R., Stacy, T., Talebian, L., Lind, E., Robin, C., Binder, M., Dzierzak, E. and Speck, N. A. (2002). Runx1 expression marks long-term repopulating hematopoietic stem cells in the midgestation mouse embryo. *Immunity* **16**, 661-672.
- Nostro, M. C., Cheng, X., Keller, G. M. and Gadue, P. (2008). Wnt, activin, and BMP signaling regulate distinct stages in the developmental pathway from embryonic stem cells to blood. *Cell Stem Cell* **2**, 60-71.
- Notta, F., Doulatov, S., Laurenti, E., Poeppl, A., Jurisica, I. and Dick, J. E. (2011). Isolation of single human hematopoietic stem cells capable of long-term multilineage engraftment. *Science* **333**, 218-221.
- Oberlin, E., Tavian, M., Blazsek, I. and Peault, B. (2002). Blood-forming potential of vascular endothelium in the human embryo. *Development* **129**, 4147-4157.
- Oberlin, E., Fleury, M., Clay, D., Petit-Cocault, L., Candelier, J.-J., Mennesson, B., Jaffredo, T. and Souyri, M. (2010). VE-cadherin expression allows identification of a new class of hematopoietic stem cells within human embryonic liver. *Blood* **116**, 4444-4455.
- Ohta, M., Sakai, T., Saga, Y., Aizawa, S. and Saito, M. (1998). Suppression of hematopoietic activity in tenascin-C-deficient mice. *Blood* **91**, 4074-4083.
- O'Rahilly, R. and Muller, F. (1987). *Developmental Stages in Human Embryos*. Washington: Carnegie Institution of Washington.
- Otersbach, K. and Dzierzak, E. (2005). The murine placenta contains hematopoietic stem cells within the vascular labyrinth region. *Dev. Cell* **8**, 377-387.
- Pereira, C.-F., Lemischka, I. R. and Moore, K. (2012). Reprogramming cell fates: insights from combinatorial approaches. *Ann. N. Y. Acad. Sci.* **1266**, 7-17.
- Pereira, C.-F., Chang, B., Qiu, X., Niu, X., Papatsenko, D., Hendry, C. E., Clark, N. R., Nomura-Kitabayashi, A., Kovacic, J. C., Ma'ayan, A. et al. (2013). Induction of a hemogenic program in mouse fibroblasts. *Cell Stem Cell* **13**, 205-218.
- Pereira, C.-F., Chang, B., Gomes, A., Bernitz, J., Papatsenko, D., Niu, X., Swiers, G., Azzoni, E., de Bruijn, M. F. T. R., Schaniel, C. et al. (2016). Hematopoietic reprogramming in vitro informs in vivo identification of hemogenic precursors to definitive hematopoietic stem cells. *Dev. Cell* **36**, 525-539.
- Pick, M., Azzola, L., Mossman, A., Stanley, E. G. and Elefanty, A. G. (2007). Differentiation of human embryonic stem cells in serum-free medium reveals distinct roles for bone morphogenetic protein 4, vascular endothelial growth factor, stem cell factor, and fibroblast growth factor 2 in hematopoiesis. *Stem Cells* **25**, 2206-2214.
- Prashad, S. L., Calvanese, V., Yao, C. Y., Kaiser, J., Wang, Y., Sasidharan, R., Crooks, G., Magnusson, M. and Mikkola, H. K. (2015). GPI-80 defines self-renewal ability in hematopoietic stem cells during human development. *Cell Stem Cell* **16**, 80-87.
- Pulecio, J., Nivet, E., Sancho-Martinez, I., Vitaloni, M., Guenachea, G., Xia, Y., Kurian, L., Dubova, I., Bueren, J., Laricchia-Robbio, L. et al. (2014). Conversion of human fibroblasts into monocyte-like progenitor cells. *Stem Cells* **32**, 2923-2938.
- Qiu, C., Olivier, E. N., Velho, M. and Bouhassira, E. E. (2008). Globin switches in yolk sac-like primitive and fetal-like definitive red blood cells produced from human embryonic stem cells. *Blood* **111**, 2400-2408.
- Rafii, S., Kloss, C. C., Butler, J. M., Ginsberg, M., Gars, E., Lis, R., Zhan, Q., Josipovic, P., Ding, B.-S., Xiang, J. et al. (2013). Human ESC-derived hemogenic endothelial cells undergo distinct waves of endothelial to hematopoietic transition. *Blood* **121**, 770-780.
- Rebel, V. I., Miller, C. L., Eaves, C. J. and Lansdorp, P. M. (1996). The repopulation potential of fetal liver hematopoietic stem cells in mice exceeds that of their liver adult bone marrow counterparts. *Blood* **87**, 3500-3507.
- Riddell, J., Gazit, R., Garrison, B. S., Guo, G., Saadatpour, A., Mandal, P. K., Ebina, W., Volchkov, P., Yuan, G.-C., Orkin, S. H. et al. (2014). Reprogramming committed murine blood cells to induced hematopoietic stem cells with defined factors. *Cell* **157**, 549-564.
- Robertson, A. L., Avagyan, S., Gansner, J. M. and Zon, L. I. (2016). Understanding the regulation of vertebrate hematopoiesis and blood disorders - big lessons from a small fish. *FEBS Lett.* **590**, 4016-4033.
- Robin, C., Bollerot, K., Mendes, S., Haak, E., Crisan, M., Cerisoli, F., Lauw, I., Kaimakis, P., Jorna, R., Vermeulen, M. et al. (2009). Human placenta is a potent



- hematopoietic niche containing hematopoietic stem and progenitor cells throughout development. *Cell Stem Cell* **5**, 385-395.
- Rosen, S. D. (2004). Ligands for L-selectin: homing, inflammation, and beyond. *Annu. Rev. Immunol.* **22**, 129-156.
- Rothstein, T. L. and Quach, T. D. (2015). The human counterpart of mouse B-1 cells. *Ann. N. Y. Acad. Sci.* **1362**, 143-152.
- Roy, A., Cowan, G., Mead, A. J., Filippi, S., Bohn, G., Chaidos, A., Tunstall, O., Chan, J. K. Y., Choolani, M., Bennett, P. et al. (2012). Perturbation of fetal liver hematopoietic stem and progenitor cell development by trisomy 21. *Proc. Natl. Acad. Sci. USA* **109**, 17579-17584.
- Rybtsov, S., Sobiesiak, M., Taoudi, S., Souilhol, C., Senserrich, J., Liakhovitskaia, A., Ivanovs, A., Frampton, J., Zhao, S. and Medvinsky, A. (2011). Hierarchical organization and early hematopoietic specification of the developing HSC lineage in the AGM region. *J. Exp. Med.* **208**, 1305-1315.
- Rybtsov, S., Batsivari, A., Bilotkach, K., Paruzina, D., Senserrich, J., Nerushev, O. and Medvinsky, A. (2014). Tracing the origin of the HSC hierarchy reveals an SCF-dependent, IL-3-independent CD43(-) embryonic precursor. *Stem Cell Rep.* **3**, 489-501.
- Sandler, V. M., Lis, R., Liu, Y., Kedem, A., James, D., Elemento, O., Butler, J. M., Scandura, J. M. and Rafii, S. (2014). Reprogramming human endothelial cells to haematopoietic cells requires vascular induction. *Nature* **511**, 312-318.
- Schier, A. F. and Shen, M. M. (2000). Nodal signalling in vertebrate development. *Nature* **403**, 385-389.
- Shultz, L. D., Schweitzer, P. A., Christianson, S. W., Gott, B., Schweitzer, I. B., Tennent, B., McKenna, S., Mobraaten, L., Rajan, T. V., Greiner, D. L. (1995). Multiple defects in innate and adaptive immunologic function in NOD/LtSz-scid mice. *J. Immunol.* **154**, 180-191.
- Shultz, L. D., Brehm, M. A., Garcia-Martinez, J. V. and Greiner, D. L. (2012). Humanized mice for immune system investigation: progress, promise and challenges. *Nat. Rev. Immunol.* **12**, 786-798.
- Silver, L. and Palis, J. (1997). Initiation of murine embryonic erythropoiesis: a spatial analysis. *Blood* **89**, 1154-1164.
- Sinka, L., Biasch, K., Khazaal, I., Peault, B. and Tavian, M. (2012). Angiotensin-converting enzyme (CD143) specifies emerging lympho-hematopoietic progenitors in the human embryo. *Blood* **119**, 3712-3723.
- Slukvin, I. I. (2016). Generating human hematopoietic stem cells in vitro -exploring endothelial to hematopoietic transition as a portal for stemness acquisition. *FEBS Lett.* **590**, 4126-4143.
- Souilhol, C., Gonneau, C., Lendinez, J. G., Batsivari, A., Rybtsov, S., Wilson, H., Morgado-Palacin, L., Hills, D., Taoudi, S., Antonchuk, J. et al. (2016). Inductive interactions mediated by interplay of asymmetric signalling underlie development of adult haematopoietic stem cells. *Nat. Commun.* **7**, 10784.
- Sturgeon, C. M., Dittadi, A., Awong, G., Kennedy, M. and Keller, G. (2014). Wnt signaling controls the specification of definitive and primitive hematopoiesis from human pluripotent stem cells. *Nat. Biotechnol.* **32**, 554-561.
- Sugimura, R., Jha, D. K., Han, A., Soria-Valles, C., da Rocha, E. L., Lu, Y.-F., Goettel, J. A., Serrao, E., Rowe, R. G., Malleshaiah, M. et al. (2017). Haematopoietic stem and progenitor cells from human pluripotent stem cells. *Nature* **545**, 432-438.
- Suna, S., Sakata, Y., Sato, H., Mizuno, H., Nakatani, D., Shimizu, M., Usami, M., Takashima, S., Takeda, H. and Hori, M. (2008). Up-regulation of cell adhesion molecule genes in human endothelial cells stimulated by lymphotoxin alpha: DNA microarray analysis. *J. Atheroscler. Thromb.* **15**, 160-165.
- Suzuki, N., Yamazaki, S., Yamaguchi, T., Okabe, M., Masaki, H., Takaki, S., Otsu, M. and Nakauchi, H. (2013). Generation of engraftable hematopoietic stem cells from induced pluripotent stem cells by way of teratoma formation. *Mol. Ther.* **21**, 1424-1431.
- Takahashi, K. and Yamanaka, S. (2006). Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* **126**, 663-676.
- Tam, P. P. L. and Behringer, R. R. (1997). Mouse gastrulation: the formation of a mammalian body plan. *Mech. Dev.* **68**, 3-25.
- Taoudi, S. and Medvinsky, A. (2007). Functional identification of the hematopoietic stem cell niche in the ventral domain of the embryonic dorsal aorta. *Proc. Natl. Acad. Sci. USA* **104**, 9399-9403.
- Taoudi, S., Morrison, A. M., Inoue, H., Gribo, R., Ure, J. and Medvinsky, A. (2005). Progressive divergence of definitive haematopoietic stem cells from the endothelial compartment does not depend on contact with the foetal liver. *Development* **132**, 4179-4191.
- Taoudi, S., Gonneau, C., Moore, K., Sheridan, J. M., Blackburn, C. C., Taylor, E. and Medvinsky, A. (2008). Extensive hematopoietic stem cell generation in the AGM region via maturation of VE-cadherin+CD45+ pre-definitive HSCs. *Cell Stem Cell* **3**, 99-108.
- Tavian, M. and Peault, B. (2005). Embryonic development of the human hematopoietic system. *Int. J. Dev. Biol.* **49**, 243-250.
- Tavian, M., Coulombel, L., Luton, D., Clemente, H. S., Dieterlen-Lievre, F. and Peault, B. (1996). Aorta-associated CD34+ hematopoietic cells in the early human embryo. *Blood* **87**, 67-72.
- Tavian, M., Hallais, M. F. and Peault, B. (1999). Emergence of intraembryonic hematopoietic precursors in the pre-liver human embryo. *Development* **126**, 793-803.
- Tavian, M., Robin, C., Coulombel, L. and Peault, B. (2001). The human embryo, but not its yolk sac, generates lympho-myeloid stem cells: mapping multipotent hematopoietic cell fate in intraembryonic mesoderm. *Immunity* **15**, 487-495.
- Taylor, E., Taoudi, S. and Medvinsky, A. (2010). Hematopoietic stem cell activity in the aorta-gonad-mesonephros region enhances after mid-day 11 of mouse development. *Int. J. Dev. Biol.* **54**, 1055-1060.
- Thambyrajah, R., Mazan, M., Patel, R., Moignard, V., Stefanska, M., Marinopoulou, E., Li, Y., Lancrin, C., Clapes, T., Möröy, T. et al. (2016). GF11 proteins orchestrate the emergence of haematopoietic stem cells through recruitment of LSD1. *Nat. Cell Biol.* **18**, 21-32.
- Timmermans, F., Velghe, I., Vanwalleghem, L., De Smedt, M., Van Coppennolle, S., Taghon, T., Moore, H. D., Leclercq, G., Langerak, A. W., Kerre, T. et al. (2009). Generation of T cells from human embryonic stem cell-derived hematopoietic zones. *J. Immunol.* **182**, 6879-6888.
- Uenishi, G., Theisen, D., Lee, J.-H., Kumar, A., Raymond, M., Vodyanik, M., Swanson, S., Stewart, R., Thomson, J. and Slukvin, I. (2014). Tenascin C promotes hematoendothelial development and T lymphoid commitment from human pluripotent stem cells in chemically defined conditions. *Stem Cell Rep.* **3**, 1073-1084.
- Van Handel, B., Prashad, S. L., Hassanzadeh-Kiabi, N., Huang, A., Magnusson, M., Atanassova, B., Chen, A., Hamalainen, E. I. and Mikkola, H. K. A. (2010). The first trimester human placenta is a site for terminal maturation of primitive erythroid cells. *Blood* **116**, 3321-3330.
- Vereide, D. T., Vickerman, V., Swanson, S. A., Chu, L.-F., McIntosh, B. E. and Thomson, J. A. (2014). An expandable, inducible hemangioblast state regulated by fibroblast growth factor. *Stem Cell Rep.* **3**, 1043-1057.
- Vervest, H. A. and Haspels, A. A. (1985). Preliminary results with the antiprogesterone compound RU-486 (mifepristone) for interruption of early pregnancy. *Fertil. Steril.* **44**, 627-632.
- Vodyanik, M. A., Thomson, J. A. and Slukvin, I. I. (2006). Leukosialin (CD43) defines hematopoietic progenitors in human embryonic stem cell differentiation cultures. *Blood* **108**, 2095-2105.
- Vodyanik, M. A., Yu, J., Zhang, X., Tian, S., Stewart, R., Thomson, J. A. and Slukvin, I. I. (2010). A mesoderm-derived precursor for mesenchymal stem and endothelial cells. *Cell Stem Cell* **7**, 718-729.
- Wahlster, L. and Daley, G. Q. (2016). Progress towards generation of human hematopoietic stem cells. *Nat. Cell Biol.* **18**, 1111-1117.
- Wang, Y. and Nakayama, N. (2009). WNT and BMP signaling are both required for hematopoietic cell development from human ES cells. *Stem Cell Res.* **3**, 113-125.
- Watt, S. M., Butler, L. H., Tavian, M., Buhring, H. J., Rappold, I., Simmons, P. J., Zannettino, A. C., Buck, D., Fuchs, A., Doyonnas, R. et al. (2000). Functionally defined CD164 epitopes are expressed on CD34(+) cells throughout ontogeny but display distinct distribution patterns in adult hematopoietic and nonhematopoietic tissues. *Blood* **95**, 3113-3124.
- Wilkinson, R. N., Pouget, C., Gering, M., Russell, A. J., Davies, S. G., Kimelman, D. and Patient, R. (2009). Hedgehog and Bmp polarize hematopoietic stem cell emergence in the zebrafish dorsal aorta. *Dev. Cell* **16**, 909-916.
- Woll, P. S., Morris, J. K., Painschab, M. S., Marcus, R. K., Kohn, A. D., Biechele, T. L., Moon, R. T. and Kaufman, D. S. (2008). Wnt signaling promotes hematoendothelial cell development from human embryonic stem cells. *Blood* **111**, 122-131.
- Yang, C.-T., French, A., Goh, P. A., Pagnamenta, A., Mettananda, S., Taylor, J., Knight, S., Nathwani, A., Roberts, D. J., Watt, S. M. et al. (2014). Human induced pluripotent stem cell derived erythroblasts can undergo definitive erythropoiesis and co-express gamma and beta globins. *Br. J. Haematol.* **166**, 435-448.
- Yokomizo, T. and Dzierzak, E. (2010). Three-dimensional cartography of hematopoietic clusters in the vasculature of whole mouse embryos. *Development* **137**, 3651-3661.
- Yoshimoto, M. (2015). The first wave of B lymphopoiesis develops independently of stem cells in the murine embryo. *Ann. N. Y. Acad. Sci.* **1362**, 16-22.
- Yoshimoto, M., Montecino-Rodriguez, E., Ferkowicz, M. J., Porayette, P., Shelley, W. C., Conway, S. J., Dorshkind, K. and Yoder, M. C. (2011). Embryonic day 9 yolk sac and intra-embryonic hemogenic endothelium independently generate a B-1 and marginal zone progenitor lacking B-2 potential. *Proc. Natl. Acad. Sci. USA* **108**, 1468-1473.
- Yoshimoto, M., Porayette, P., Glosson, N. L., Conway, S. J., Carlesso, N., Cardoso, A. A., Kaplan, M. H. and Yoder, M. C. (2012). Autonomous murine T-cell progenitor production in the extra-embryonic yolk sac before HSC emergence. *Blood* **119**, 5706-5714.
- Yu, P., Pan, G., Yu, J. and Thomson, J. A. (2011). FGF2 sustains NANOG and switches the outcome of BMP4-induced human embryonic stem cell differentiation. *Cell Stem Cell* **8**, 326-334.
- Yu, Q. C., Hirst, C. E., Costa, M., Ng, E. S., Schiesser, J. V., Gertow, K., Stanley, E. G. and Elefanty, A. G. (2012). APELIN promotes hematopoiesis from human embryonic stem cells. *Blood* **119**, 6243-6254.
- Zhou, F., Li, X., Wang, W., Zhu, P., Zhou, J., He, W., Ding, M., Xiong, F., Zheng, X., Li, Z. et al. (2016). Tracing haematopoietic stem cell formation at single-cell resolution. *Nature* **533**, 487-492.
- Zovein, A. C., Hofmann, J. J., Lynch, M., French, W. J., Turlo, K. A., Yang, Y., Becker, M. S., Zanetta, L., Dejana, E., Gasson, J. C. et al. (2008). Fate tracing reveals the endothelial origin of hematopoietic stem cells. *Cell Stem Cell* **3**, 625-636.